

PCT COOPERATION TREATY

PCT

NOTIFICATION OF ELECTION

(PCT Rule 61.2)

From the INTERNATIONAL BUREAU

To:

Commissioner
US Department of Commerce
United States Patent and Trademark
Office, PCT
2011 South Clark Place Room
CP2/5C24
Arlington, VA 22202
ETATS-UNIS D'AMERIQUE

in its capacity as elected Office

Date of mailing (day/month/year) 17 August 2001 (17.08.01)	
International application No. PCT/SE01/00248	Applicant's or agent's file reference Z70663-1 WO
International filing date (day/month/year) 07 February 2001 (07.02.01)	Priority date (day/month/year) 12 February 2000 (12.02.00)
Applicant BAXTER, Andrew et al	

1. The designated Office is hereby notified of its election made:

☒ in the demand filed with the International Preliminary Examining Authority on:

23 July 2001 (23.07.01)

☐ in a notice effecting later election filed with the International Bureau on:

2. The election
- ☒
- was

☐ was not

made before the expiration of 19 months from the priority date or, where Rule 32 applies, within the time limit under Rule 32.2(b).

The International Bureau of WIPO 34, chemin des Colombettes 1211 Geneva 20, Switzerland Facsimile No.: (41-22) 740.14.35	Authorized officer Charlotte ENGER Telephone No.: (41-22) 338.83.38
---	---

PATENT COOPERATION TREATY

PCT

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Article 36 and Rule 70)

8
REC'D 29 MAY 2002

WIPO PCT

9/868884

Applicant's or agent's file reference Z70663-1 WO	FOR FURTHER ACTION See Notification of Transmittal of International Preliminary Examination Report (Form PCT/IPEA/416)	
International application No. PCT/SE01/00248	International filing date (day/month/year) 07.02.2001	Priority date (day/month/year) 12.02.2000
International Patent Classification (IPC) or national classification and IPC7 C07D 333/38, 409/04, 413/04, 417/04, 263/48, A61K 31/381, 31/395, A61P 29/00, 11/06, 19/02		
Applicant AstraZeneca AB et al		NOV 01 2002

RECEIVED

TECH CENTER 1600/2900

1. This international preliminary examination report has been prepared by this International Preliminary Examining Authority and is transmitted to the applicant according to Article 36.

2. This REPORT consists of a total of 5 sheets, including this cover sheet.

☐ This report is also accompanied by ANNEXES, i.e., sheets of the description, claims and/or drawings which have been amended and are the basis for this report and/or sheets containing rectifications made before this Authority (see Rule 70.16 and Section 607 of the Administrative Instructions under the PCT).

These annexes consist of a total of _____ sheets.

3. This report contains indications relating to the following items:

- I ☒ Basis of the report
- II ☐ Priority
- III ☒ Non-establishment of opinion with regard to novelty, inventive step and industrial applicability
- IV ☐ Lack of unity of invention
- V ☒ Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement
- VI ☐ Certain documents cited
- VII ☐ Certain defects in the international application
- VIII ☐ Certain observations on the international application

Date of submission of the demand 23.07.2001	Date of completion of this report 13.05.2002
Name and mailing address of the IPEA/SE Patent- och registreringsverket Box 5055 S-102 42 STOCKHOLM Facsimile No. 08-667 72 88	Authorized officer Nebil Gecer/EÖ Telephone No. 08-782 25 00

Form PCT/IPEA/409 (cover sheet) (January 1998)

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No.

PCT/SE01/00248

I. Basis of the report

1. With regard to the elements of the international application:*

☒ the international application as originally filed☐ the description:

pages _____, as originally filed

pages _____, filed with the demand

pages _____, filed with the letter of _____

☐ the claims:

pages _____, as originally filed

pages _____, as amended (together with any statement) under article 19

pages _____, filed with the demand

pages _____, filed with the letter of _____

☐ the drawings:

pages _____, as originally filed

pages _____, filed with the demand

pages _____, filed with the letter of _____

☐ the sequence listing part of the description:

pages _____, as originally filed

pages _____, filed with the demand

pages _____, filed with the letter of _____

2. With regard to the language, all the elements marked above were available or furnished to this Authority in the language in which the international application was filed, unless otherwise indicated under this item.

These elements were available or furnished to this Authority in the following language _____ which is:

☐ the language of a translation furnished for the purposes of international search (under Rule 23.1(b)).☐ the language of publication of the international application (under Rule 48.3(b)).☐ the language of the translation furnished for the purposes of international preliminary examination (under Rules 55.2 and/or 55.3).

3. With regard to any nucleotide and/or amino acid sequence disclosed in the international application, the international preliminary examination was carried out on the basis of the sequence listing:

☐ contained in the international application in written form.☐ filed together with the international application in computer readable form.☐ furnished subsequently to this Authority in written form.☐ furnished subsequently to this Authority in computer readable form.☐ The statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.☐ The statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished.4. ☐ The amendments have resulted in the cancellation of:☐ the description, pages _____☐ the claims, Nos. _____☐ the drawings, sheet/fig _____5. ☐ This report has been established as if (some of) the amendments had not been made, since they have been considered to go beyond the disclosure as filed, as indicated in the Supplemental Box (Rule 70.2 (c)).**

* Replacement sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this report as "originally filed" and are annexed to this report since they do not contain amendments (Rules 70.16 and 70.17).

** Any replacement sheet containing such amendments must be referred to under item I and annexed to this report.

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No.

PCT/SE01/00248

III. Non-establishment of opinion with regard to novelty, inventive step and industrial applicability

1. The questions whether the claimed invention appears to be novel, to involve an inventive step (to be non obvious), or to be industrially applicable have not been examined in respect of:

☐ the entire international application,

☒ claims Nos. 20-25

because:

☒ the said international application, or the said claims Nos. 20-25
relate to the following subject matter which does not require an international preliminary examination (*specify*):

See PCT Rule 67.1.(iv).: Methods for treatment of the human or animal body by surgery or therapy, as well as diagnostic methods.

☐ the description, claims or drawings (*indicate particular elements below*) or said claims Nos. _____
are so unclear that no meaningful opinion could be formed (*specify*):

☐ the claims, or said claims Nos. _____ are so inadequately supported
by the description that no meaningful opinion could be formed.

☐ no international search report has been established for said claims Nos. _____

2. A meaningful international preliminary examination cannot be carried out due to the failure of the nucleotide and/or amino acid sequence listing to comply with the standard provided for in Annex C of the Administrative Instructions:

☐ the written form has not been furnished or does not comply with the standard.

☐ the computer readable form has not been furnished or does not comply with the standard.

PCT/SE01/00248

1. Statement

YES

NO

YES

NO

YES

NO

.../...

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No.

PCT/SE01/00248

Supplemental Box

(To be used when the space in any of the preceding boxes is not sufficient)

Continuation of: V

If an inventive step is shown to exist, it must be shown that the unexpected effect is due to a distinguishing feature, which exists for the whole scope of protection.

D4 relates to structurally more remote pharmaceutically active 3-pyrazolyl derivatives and D5 reveals structurally related substituted thienyl compounds and their use as growth promoters for animals. D4 and D5 only disclose the general state of the art and are not considered to be of particular relevance.

PCT

REQUEST

The undersigned requests that the present international application be processed according to the Patent Cooperation Treaty.

Receiving Office use only

International Application No.

International Filing Date

Name of receiving Office and "PCT International Application"

Applicant's or agent's file reference
(if desired) (12 characters maximum) Z70663-1 WO

Box No. I TITLE OF INVENTION
NOVEL COMPOUNDS

Box No. II APPLICANT

Name and address: (Family name followed by given name; for a legal entity, full official designation. The address must include postal code and name of country. The country of the address indicated in this Box is the applicant's State (that is, country) of residence if no State of residence is indicated below.)

AstraZeneca AB
S-151 85 Södertälje
Sweden

☐ This person is also inventor.

Telephone No.

+46 8 553 260 00

Facsimile No.

+46 8 553 288 20

Teleprinter No.

State (that is, country) of nationality:
SE

State (that is, country) of residence:
SE

This person is applicant for the purposes of: ☐ all designated States ☒ all designated States except the United States of America ☐ the United States of America only ☐ the States indicated in the Supplemental Box

Box No. III FURTHER APPLICANT(S) AND/OR (FURTHER) INVENTOR(S)

Name and address: (Family name followed by given name; for a legal entity, full official designation. The address must include postal code and name of country. The country of the address indicated in this Box is the applicant's State (that is, country) of residence if no State of residence is indicated below.)

BAXTER, Andrew
AstraZeneca R&D Charnwood
Bakewell Road
Loughborough, Leics. LE11 5RH
United Kingdom

This person is:

☐ applicant only

☒ applicant and inventor

☐ inventor only (If this check-box is marked, do not fill in below.)

State (that is, country) of nationality:
GB

State (that is, country) of residence:
GB

This person is applicant for the purposes of: ☐ all designated States ☐ all designated States except the United States of America ☒ the United States of America only ☐ the States indicated in the Supplemental Box

☒ Further applicants and/or (further) inventors are indicated on a continuation sheet.

Box No. IV AGENT OR COMMON REPRESENTATIVE; OR ADDRESS FOR CORRESPONDENCE

The person identified below is hereby/has been appointed to act on behalf of the applicant(s) before the competent International Authorities as: ☒ agent ☐ common representative

Name and address: (Family name followed by given name; for a legal entity, full official designation. The address must include postal code and name of country.)

Global Intellectual Property
AstraZeneca AB
S-151 85 Södertälje
Sweden

Telephone No.

+46 8 553 260 00

Facsimile No.

+46 8 553 288 20

Teleprinter No.

☐ Address for correspondence: Mark this check-box where no agent or common representative is/has been appointed and the space above is used instead to indicate a special address to which correspondence should be sent.

Continuation of Box No. II. FURTHER APPLICANT(S) AND/OR (FURTHER) INVENTOR(S)	
<i>If none of the following sub-boxes is used, this sheet should not be included in the request.</i>	
<p>Name and address: <i>(Family name followed by given name; for a legal entity, full official designation. The address must include postal code and name of country. The country of the address indicated in this Box is the applicant's State (that is, country) of residence if no State of residence is indicated below.)</i></p> <p>BROUGH, Stephen AstraZeneca R&D Charnwood Bakewell Road Loughborough, Leics. LE11 5RH United Kingdom</p>	<p>This person is:</p> <p><input type="checkbox"/> applicant only</p> <p><input checked="" type="checkbox"/> applicant and inventor</p> <p><input type="checkbox"/> inventor only <i>(If this check-box is marked, do not fill in below.)</i></p>
State <i>(that is, country)</i> of nationality: GB	State <i>(that is, country)</i> of residence: GB
<p>This person is applicant for the purposes of: <input type="checkbox"/> all designated States <input type="checkbox"/> all designated States except the United States of America <input checked="" type="checkbox"/> the United States of America only <input type="checkbox"/> the States indicated in the Supplemental Box</p>	
<p>Name and address: <i>(Family name followed by given name; for a legal entity, full official designation. The address must include postal code and name of country. The country of the address indicated in this Box is the applicant's State (that is, country) of residence if no State of residence is indicated below.)</i></p> <p>FAULL, Alan AstraZeneca R&D Alderley Mereseide, Alderley Park, Macclesfield, Cheshire SK10 4TG England</p>	<p>This person is:</p> <p><input type="checkbox"/> applicant only</p> <p><input checked="" type="checkbox"/> applicant and inventor</p> <p><input type="checkbox"/> inventor only <i>(If this check-box is marked, do not fill in below.)</i></p>
State <i>(that is, country)</i> of nationality: GB	State <i>(that is, country)</i> of residence: GB
<p>This person is applicant for the purposes of: <input type="checkbox"/> all designated States <input type="checkbox"/> all designated States except the United States of America <input checked="" type="checkbox"/> the United States of America only <input type="checkbox"/> the States indicated in the Supplemental Box</p>	
<p>Name and address: <i>(Family name followed by given name; for a legal entity, full official designation. The address must include postal code and name of country. The country of the address indicated in this Box is the applicant's State (that is, country) of residence if no State of residence is indicated below.)</i></p> <p>JOHNSTONE, Craig AstraZeneca R&D Alderley Mereseide, Alderley Park, Macclesfield, Cheshire SK10 4TG England</p>	<p>This person is:</p> <p><input type="checkbox"/> applicant only</p> <p><input checked="" type="checkbox"/> applicant and inventor</p> <p><input type="checkbox"/> inventor only <i>(If this check-box is marked, do not fill in below.)</i></p>
State <i>(that is, country)</i> of nationality: GB	State <i>(that is, country)</i> of residence: GB
<p>This person is applicant for the purposes of: <input type="checkbox"/> all designated States <input type="checkbox"/> all designated States except the United States of America <input checked="" type="checkbox"/> the United States of America only <input type="checkbox"/> the States indicated in the Supplemental Box</p>	
<p>Name and address: <i>(Family name followed by given name; for a legal entity, full official designation. The address must include postal code and name of country. The country of the address indicated in this Box is the applicant's State (that is, country) of residence if no State of residence is indicated below.)</i></p> <p>MCINALLY, Thomas AstraZeneca R&D Charnwood Bakewell Road Loughborough, Leics. LE11 5RH United Kingdom</p>	<p>This person is:</p> <p><input type="checkbox"/> applicant only</p> <p><input checked="" type="checkbox"/> applicant and inventor</p> <p><input type="checkbox"/> inventor only <i>(If this check-box is marked, do not fill in below.)</i></p>
State <i>(that is, country)</i> of nationality: GB	State <i>(that is, country)</i> of residence: GB
<p>This person is applicant for the purposes of: <input type="checkbox"/> all designated States <input type="checkbox"/> all designated States except the United States of America <input checked="" type="checkbox"/> the United States of America only <input type="checkbox"/> the States indicated in the Supplemental Box</p>	
<p><input type="checkbox"/> Further applicants and/or (further) inventors are indicated on another continuation sheet.</p>	

Box No. V DESIGNATION OF STATES

The following designations are hereby made under Rule 4.9(a) (mark the applicable check-boxes; at least one must be marked):

Regional Patent

- ☒ AP ARIPO Patent: GH Ghana, GM Gambia, KE Kenya, LS Lesotho, MW Malawi, MZ Mozambique, SD Sudan, SL Sierra Leone, SZ Swaziland, TZ United Republic of Tanzania, UG Uganda, ZW Zimbabwe, and any other State which is a Contracting State of the Harare Protocol and of the PCT
- ☒ EA Eurasian Patent: AM Armenia, AZ Azerbaijan, BY Belarus, KG Kyrgyzstan, KZ Kazakhstan, MD Republic of Moldova, RU Russian Federation, TJ Tajikistan, TM Turkmenistan, and any other State which is a Contracting State of the Eurasian Patent Convention and of the PCT
- ☒ EP European Patent: AT Austria, BE Belgium, CH and LI Switzerland and Liechtenstein, CY Cyprus, DE Germany, DK Denmark, ES Spain, FI Finland, FR France, GB United Kingdom, GR Greece, IE Ireland, IT Italy, LU Luxembourg, MC Monaco, NL Netherlands, PT Portugal, SE Sweden, TR Turkey, and any other State which is a Contracting State of the European Patent Convention and of the PCT
- ☒ OA OAPI Patent: BF Burkina Faso, BJ Benin, CF Central African Republic, CG Congo, CI Côte d'Ivoire, CM Cameroon, GA Gabon, GN Guinea, GW Guinea-Bissau, ML Mali, MR Mauritania, NE Niger, SN Senegal, TD Chad, TG Togo, and any other State which is a member State of OAPI and a Contracting State of the PCT (if other kind of protection or treatment desired, specify on dotted line)


National Patent (if other kind of protection or treatment desired, specify on dotted line):

- | | |
|--|--|
| <input checked="" type="checkbox"/> AE United Arab Emirates | <input checked="" type="checkbox"/> LC Saint Lucia |
| <input checked="" type="checkbox"/> AG Antigua and Barbuda | <input checked="" type="checkbox"/> LK Sri Lanka |
| <input checked="" type="checkbox"/> AL Albania | <input checked="" type="checkbox"/> LR Liberia |
| <input checked="" type="checkbox"/> AM Armenia | <input checked="" type="checkbox"/> LS Lesotho |
| <input checked="" type="checkbox"/> AT Austria | <input checked="" type="checkbox"/> LT Lithuania |
| <input checked="" type="checkbox"/> AU Australia | <input checked="" type="checkbox"/> LU Luxembourg |
| <input checked="" type="checkbox"/> AZ Azerbaijan | <input checked="" type="checkbox"/> LV Latvia |
| <input checked="" type="checkbox"/> BA Bosnia and Herzegovina | <input checked="" type="checkbox"/> MA Morocco |
| <input checked="" type="checkbox"/> BB Barbados | <input checked="" type="checkbox"/> MD Republic of Moldova |
| <input checked="" type="checkbox"/> BG Bulgaria | <input checked="" type="checkbox"/> MG Madagascar |
| <input checked="" type="checkbox"/> BR Brazil | <input checked="" type="checkbox"/> MK The former Yugoslav Republic of Macedonia |
| <input checked="" type="checkbox"/> BY Belarus | <input checked="" type="checkbox"/> MN Mongolia |
| <input checked="" type="checkbox"/> BZ Belize | <input checked="" type="checkbox"/> MW Malawi |
| <input checked="" type="checkbox"/> CA Canada | <input checked="" type="checkbox"/> MX Mexico |
| <input checked="" type="checkbox"/> CH and LI Switzerland and Liechtenstein | <input checked="" type="checkbox"/> MZ Mozambique |
| <input checked="" type="checkbox"/> CN China | <input checked="" type="checkbox"/> NO Norway |
| <input checked="" type="checkbox"/> CR Costa Rica | <input checked="" type="checkbox"/> NZ New Zealand |
| <input checked="" type="checkbox"/> CU Cuba | <input checked="" type="checkbox"/> PL Poland |
| <input checked="" type="checkbox"/> CZ Czech Republic | <input checked="" type="checkbox"/> PT Portugal |
| <input checked="" type="checkbox"/> DE Germany | <input checked="" type="checkbox"/> RO Romania |
| <input checked="" type="checkbox"/> DK Denmark | <input checked="" type="checkbox"/> RU Russian Federation |
| <input checked="" type="checkbox"/> DM Dominica | <input checked="" type="checkbox"/> SD Sudan |
| <input checked="" type="checkbox"/> DZ Algeria | <input checked="" type="checkbox"/> SE Sweden |
| <input checked="" type="checkbox"/> EE Estonia | <input checked="" type="checkbox"/> SG Singapore |
| <input checked="" type="checkbox"/> ES Spain | <input checked="" type="checkbox"/> SI Slovenia |
| <input checked="" type="checkbox"/> FI Finland | <input checked="" type="checkbox"/> SK Slovakia |
| <input checked="" type="checkbox"/> GB United Kingdom | <input checked="" type="checkbox"/> SL Sierra Leone |
| <input checked="" type="checkbox"/> GD Grenada | <input checked="" type="checkbox"/> TJ Tajikistan |
| <input checked="" type="checkbox"/> GE Georgia | <input checked="" type="checkbox"/> TM Turkmenistan |
| <input checked="" type="checkbox"/> GH Ghana | <input checked="" type="checkbox"/> TR Turkey |
| <input checked="" type="checkbox"/> GM Gambia | <input checked="" type="checkbox"/> TT Trinidad and Tobago |
| <input checked="" type="checkbox"/> HR Croatia | <input checked="" type="checkbox"/> TZ United Republic of Tanzania |
| <input checked="" type="checkbox"/> HU Hungary | <input checked="" type="checkbox"/> UA Ukraine |
| <input checked="" type="checkbox"/> ID Indonesia | <input checked="" type="checkbox"/> UG Uganda |
| <input checked="" type="checkbox"/> IL Israel | <input checked="" type="checkbox"/> US United States of America |
| <input checked="" type="checkbox"/> IN India | <input checked="" type="checkbox"/> UZ Uzbekistan |
| <input checked="" type="checkbox"/> IS Iceland | <input checked="" type="checkbox"/> VN Viet Nam |
| <input checked="" type="checkbox"/> JP Japan | <input checked="" type="checkbox"/> YU Yugoslavia |
| <input checked="" type="checkbox"/> KE Kenya | <input checked="" type="checkbox"/> ZA South Africa |
| <input checked="" type="checkbox"/> KG Kyrgyzstan | <input checked="" type="checkbox"/> ZW Zimbabwe |
| <input checked="" type="checkbox"/> KP Democratic People's Republic of Korea | |
| <input checked="" type="checkbox"/> KR Republic of Korea | |
| <input checked="" type="checkbox"/> KZ Kazakhstan | |

Check-box reserved for designating States which have become party to the PCT after issuance of this sheet:



Precautionary Designation Statement: In addition to the designations made above, the applicant also makes under Rule 4.9(b) all other designations which would be permitted under the PCT except any designation(s) indicated in the Supplemental Box as being excluded from the scope of this statement. The applicant declares that those additional designations are subject to confirmation and that any designation which is not confirmed before the expiration of 15 months from the priority date is to be regarded as withdrawn by the applicant at the expiration of that time limit. (Confirmation (including fees) must reach the receiving Office within the 15-month time limit.)

Box No. VI PRIORITY CLAIM					<input type="checkbox"/> Further priority claims are indicated in the Supplemental Box.
Filing date of earlier application (day/month/year)	Number of earlier application	Where earlier application is:			
		national application: country	regional application:* regional Office	international application: receiving Office	
item (1) 12.02.2000 (12 February 2000)	0003154.2	Great Britain (GB)			
item (2)					
item (3)					
<input type="checkbox"/> The receiving Office is requested to prepare and transmit to the International Bureau a certified copy of the earlier application(s) (only if the earlier application was filed with the Office which for the purposes of the present international application is the receiving Office) identified above as item(s):					
* Where the earlier application is an ARIPO application, it is mandatory to indicate in the Supplemental Box at least one country party to the Paris Convention for the Protection of Industrial Property for which that earlier application was filed (Rule 4.10(b)(ii)). See Supplemental Box.					
Box No. VII INTERNATIONAL SEARCHING AUTHORITY					
Choice of International Searching Authority (ISA) (if two or more International Searching Authorities are competent to carry out the international search, indicate the Authority chosen; the two-letter code may be used):		Request to use results of earlier search; reference to that search (if an earlier search has been carried out by or requested from the International Searching Authority):			
ISA / SE		Date (day/month/year)	Number	Country (or regional Office)	
Box No. VIII CHECK LIST; LANGUAGE OF FILING					
This international application contains the following number of sheets: request : 4 description (excluding sequence listing part) : 68 claims : 11 abstract : 1 drawings : sequence listing part of description : Total number of sheets : 84		This international application is accompanied by the item(s) marked below: 1. <input checked="" type="checkbox"/> fee calculation sheet 2. <input type="checkbox"/> separate signed power of attorney 3. <input checked="" type="checkbox"/> copy of general power of attorney; reference number, if any: GF53/2001/AD 4. <input type="checkbox"/> statement explaining lack of signature 5. <input type="checkbox"/> priority document(s) identified in Box No. VI as item(s): 6. <input type="checkbox"/> translation of international application into (language): 7. <input type="checkbox"/> separate indications concerning deposited microorganism or other biological material 8. <input type="checkbox"/> nucleotide and/or amino acid sequence listing in computer readable form 9. <input type="checkbox"/> other (specify):			
Figure of the drawings which should accompany the abstract:		Language of filing of the international application: English			
Box No. IX SIGNATURE OF APPLICANT OR AGENT					
Next to each signature, indicate the name of the person signing and the capacity in which the person signs (if such capacity is not obvious from reading the request).					
Södertälje, 7 February 2001  Gunilla Fredriksson Global Intellectual Property, AstraZeneca AB					

For receiving Office use only	
1. Date of actual receipt of the purported international application: 3. Corrected date of actual receipt due to later but timely received papers or drawings completing the purported international application: 4. Date of timely receipt of the required corrections under PCT Article 11(2): 5. International Searching Authority (if two or more are competent): ISA /	2. Drawings: <input type="checkbox"/> received: <input type="checkbox"/> not received: 6. <input type="checkbox"/> Transmittal of search copy delayed until search fee is paid.

For International Bureau use only
Date of receipt of the record copy by the International Bureau:

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
16 August 2001 (16.08.2001)

PCT

(10) International Publication Number
WO 01/58890 A1

(51) International Patent Classification⁷: C07D 333/38,
409/04, 413/04, 417/04, 263/48, A61K 31/381, 31/395,
A61P 29/00, 11/06, 19/02

Thomas [GB/GB]; AstraZeneca R & D Charnwood,
Bakewell Road, Loughborough, Leics. LE11 5RH (GB).

(21) International Application Number: PCT/SE01/00248

(74) Agent: GLOBAL INTELLECTUAL PROPERTY; As-
traZeneca AB, S-151 85 Södertälje (SE).

(22) International Filing Date: 7 February 2001 (07.02.2001)

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0003154.2 12 February 2000 (12.02.2000) GB

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and


(75) Inventors/Applicants (*for US only*): BAXTER, An-
drew [GB/GB]; AstraZeneca R & D Charnwood,
Bakewell Road, Loughborough, Leics. LE11 5RH (GB).
BROUGH, Stephen [GB/GB]; AstraZeneca R & D
Charnwood, Bakewell Road, Loughborough, Leics. LE11
5RH (GB). FAULL, Alan [GB/GB]; AstraZeneca R & D
Alderley, Mereside, Alderley Park, Macclesfield, Cheshire
SK10 4TG (GB). JOHNSTONE, Craig [GB/GB]; As-
traZeneca R & D Alderley, Mereside, Alderley Park,
Macclesfield, Cheshire SK10 4TG (GB). MCINALLY,

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: HETEROAROMATIC CARBOXAMIDE DERIVATIVES AND THEIR USE AS INHIBITORS OF THE ENZYME
IKK-2



(I)

(57) Abstract: The invention relates to heteroaromatic carboxam-
ides of formula (I), wherein A, R¹, R² and X are as defined in the
specification, processes and intermediates used in their prepara-
tion, pharmaceutical compositions containing them and their use
in therapy.

WO 01/58890 A1

HETEROAROMATIC CARBOXAMIDE DERIVATIVES AND THEIR USE AS INHIBITORS OF THE ENZYME IKK-2

Field of the Invention

- 5 The present invention relates to heteroaromatic carboxamide derivatives, processes and intermediates used in their preparation, pharmaceutical compositions containing them and their use in therapy.

Background of the Invention

- 10 The NF- κ B (nuclear factor κ B) family is composed of homo- and heterodimers of the Rel family of transcription factors. A key role of these transcription factors is to induce and co-ordinate the expression of a broad spectrum of pro-inflammatory genes including cytokines, chemokines, interferons, MHC proteins, growth factors and cell adhesion
15 molecules (for reviews see Verma et. al., Genes Dev. 9:2723-35, 1995; Siebenlist et. al., Ann. Rev. Cell. Biol. 10:405-455, 1994; Bauerle and Henkel, Ann. Rev. Immunol., 12:141-179, 1994; Barnes and Karin, New Engl. J. Med., 336:1066-1071, 1997).

- The most commonly found Rel family dimer complex is composed of p50 NF κ B and p65
20 RelA (Baeuerle and Baltimore, Cell 53:211-217, 1988; Baeuerle and Baltimore, Genes Dev. 3:1689-1698, 1989). Under resting conditions NF- κ B dimers are retained in the cytoplasm by a member of the I κ B family of inhibitory proteins (Beg et. al., Genes Dev., 7:2064-2070, 1993; Gilmore and Morin, Trends Genet. 9:427-433, 1993; Haskil et. al., Cell 65:1281-1289, 1991). However, upon cell activation by a variety of cytokines or other
25 external stimuli, I κ B proteins become phosphorylated on two critical serine residues (Traenckner et. al., EMBO J., 14:2876, 1995) and are then targeted for ubiquitination and proteosome-mediated degradation (Chen, Z.J. et. al., Genes and Dev. 9:1586-1597, 1995; Scherer, D.C. et. al., Proc. Natl. Acad. Sci. USA 92:11259-11263, 1996; Alkalay, I. et. al., Proc. Natl. Acad. Sci. USA 92:10599-10603, 1995). The released NF- κ B is then able to
30 translocate to the nucleus and activate gene transcription (Beg et.al., Genes Dev., 6:1899-1913, 1992).

A wide range of external stimuli have been shown to be capable of activating NF- κ B (Baeuerle, P.A., and Baichwal, V.R., *Adv. Immunol.*, 65:111-136, 1997). Although the majority of NF- κ B activators result in I κ B phosphorylation, it is clear that multiple
5 pathways lead to this key event. Receptor-mediated NF- κ B activation relies upon specific interactions between the receptor and adapter/signalling molecules (for example, TRADD, RIP, TRAF, MyD88) and associated kinases (IRAK, NIK) (Song et. al., *Proc. Natl. Acad. Sci. USA* 94:9792-9796, 1997; Natoli et. al., *JBC* 272:26079-26082, 1997). Environmental stresses such as UV light and γ -radiation appear to stimulate NF- κ B via alternative, less
10 defined, mechanisms.

Recent publications have partially elucidated the NF- κ B activation. This work has identified three key enzymes which regulate specific I κ B/NF- κ B interactions: NF- κ B inducing kinase (NIK) (Boldin et. al., *Cell* 85:803-815, 1996), I κ B kinase-1 (IKK-1)
15 (Didonato et. al., *Nature* 388:548, 1997; Regnier et. al., *Cell* 90:373 1997) and I κ B kinase-2 (IKK-2) (Woronicz et. al., *Science* 278:866, 1997; Zandi et. al., *Cell* 91:243, 1997).

NIK appears to represent a common mediator of NF- κ B signalling cascades triggered by tumour necrosis factor and interleukin-1, and is a potent inducer of I κ B phosphorylation.
20 However NIK is unable to phosphorylate I κ B directly.

IKK-1 and IKK-2 are thought to lie immediately downstream of NIK and are capable of directly phosphorylating all three I κ B sub-types. IKK-1 and IKK-2 are 52% identical at the amino acid level but appear to have similar substrate specificities; however, enzyme
25 activities appear to be different: IKK-2 is several-fold more potent than IKK-1. Expression data, coupled with mutagenesis studies, suggest that IKK-1 and IKK-2 are capable of forming homo- and heterodimers through their C-terminal leucine zipper motifs, with the heterodimeric form being preferred (Mercurio et. al., *Mol. Cell Biol.*, 19:1526, 1999; Zandi et. al., *Science*; 281:1360, 1998; Lee et. al, *Proc. Natl. Acad. Sci. USA* 95:9319, 1998).

NIK, IKK-1 and IKK-2 are all serine/threonine kinases. Recent data has shown that tyrosine kinases also play a role in regulating the activation of NF- κ B. A number of groups have shown that TNF- α induced NF- κ B activation can be regulated by protein tyrosine phosphatases (PTPs) and tyrosine kinases (Amer et. al., JBC 273:29417-29423, 1998; Hu et. al., JBC 273:33561-33565, 1998; Kaekawa et. al., Biochem. J. 337:179-184, 1999; Singh et. al., JBC 271 31049-31054, 1996). The mechanism of action of these enzymes appears to be in regulating the phosphorylation status of I κ B. For example, PTP1B and an unidentified tyrosine kinase appear to directly control the phosphorylation of a lysine residue (K42) on I κ B- α , which in turn has a critical influence on the accessibility of the adjacent serine residues as targets for phosphorylation by IKK.

Several groups have shown that IKK-1 and IKK-2 form part of a 'signalosome' structure in association with additional proteins including IKAP (Cohen et. al., Nature 395:292-296, 1998; Rothwarf et. al., Nature 395:297-300, 1998), MEKK-1, putative MAP kinase phosphatase (Lee et. al., Proc. Natl. Acad. Sci. USA 95:9319-9324, 1998), as well as NIK and I κ B. Data is now emerging to suggest that although both IKK-1 and IKK-2 associate with NIK, they are differentially activated, and therefore might represent an important integration point for the spectrum of signals that activate NF- κ B. Importantly, MEKK-1 (one of the components of the putative signalosome and a target for UV light, LPS induced signalling molecules and small GTPases) has been found to activate IKK-2 but not IKK-1. Similarly, NIK phosphorylation of IKK-1 results in a dramatic increase in IKK-1 activity but only a small effect on IKK-2 (for review, see Mercurio, F., and Manning, A.M., Current Opinion in Cell Biology, 11:226-232, 1999).

Inhibition of NF- κ B activation is likely to be of broad utility in the treatment of inflammatory disease.

WO 98/02430 and EP 853 083 disclose various 4-pyridyl derivatives, and EP 908 456 discloses various 3-pyrazolyl derivatives.


DE 19725450 discloses various 3-pyridinyl and 5-pyrimidyl derivatives.

WO 99/46244, WO 98/54116 and EP 202 538 disclose a series of substituted thienyl compounds said to possess biological activity.

5

Disclosure of the Invention

According to the present invention, there is provided a compound of formula (I)



10

in which:

A represents a 5-membered heteroaromatic ring containing one or two heteroatoms
15 selected independently from oxygen, nitrogen or sulfur;

R¹ represents a phenyl group or a 5- to 7-membered heteroaromatic ring containing one to
three heteroatoms selected independently from oxygen, nitrogen or sulfur; said phenyl or
heteroaromatic ring being optionally substituted by one or more substituents selected
20 independently from halogen, cyano, nitro, -NR³R⁴, -CONR⁵R⁶, -COOR⁷, -NR⁸COR⁹,
-SR¹⁰, -S(O)_mR¹⁰, -SO₂NR⁵R⁶, -NR⁸SO₂R¹⁰, C₁-C₆ alkyl, trifluoromethyl, -(CH₂)_nR¹¹,
-O(CH₂)_nR¹¹ or -OR¹²;

R² represents hydrogen, halogen, cyano, nitro, -NR¹³R¹⁴, -CONR¹⁵R¹⁶, -COOR¹⁷,

$-\text{NR}^{18}\text{COR}^{19}$, $-\text{S}(\text{O})_m\text{R}^{20}$, $-\text{SO}_2\text{NR}^{15}\text{R}^{16}$, $-\text{NR}^{18}\text{SO}_2\text{R}^{20}$, $\text{C}_1\text{-C}_2$ alkyl, trifluoromethyl, $\text{C}_2\text{-C}_3$ alkenyl, $\text{C}_2\text{-C}_3$ alkynyl, trifluoromethoxy, $\text{C}_1\text{-C}_2$ alkoxy or $\text{C}_1\text{-C}_2$ alkanoyl;

X represents oxygen or sulphur;

5

each of R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , R^9 , R^{10} and R^{12} independently represent a hydrogen atom or $\text{C}_1\text{-C}_6$ alkyl;

R^{11} represents $\text{NR}^{21}\text{R}^{22}$ where R^{21} and R^{22} are independently hydrogen or $\text{C}_1\text{-C}_6$ alkyl optionally substituted by $\text{C}_1\text{-C}_4$ alkoxy; or R^{21} and R^{22} together with the nitrogen atom to which they are attached form a 5- or 6-membered saturated ring optionally containing a further O, S or NR^{23} group where R^{23} is hydrogen or $\text{C}_1\text{-C}_6$ alkyl; or R^{11} represents OR^{24} where R^{24} represents $\text{C}_1\text{-C}_6$ alkyl;

15 each of R^{13} , R^{14} , R^{15} , R^{16} , R^{17} , R^{18} , R^{19} and R^{20} independently represent a hydrogen atom or $\text{C}_1\text{-C}_2$ alkyl;

m represents an integer 0, 1 or 2;

20 n represents an integer 2, 3 or 4;

and optical isomers, racemates and tautomers thereof and pharmaceutically acceptable salts or solvates thereof:

25 provided that:

when A represents thiophene, furan or pyrrole, then R^1 is not 4-pyridinyl or 3-pyrazolyl; and

when A represents oxazole, thiazole or imidazole, then R^1 is not 3-pyridinyl or 5-pyrimidyl.

30

Certain compounds of formula (I) are capable of existing in stereoisomeric forms. It will be understood that the invention encompasses all geometric and optical isomers of the compounds of formula (I) and mixtures thereof including racemates. Tautomers and mixtures thereof also form an aspect of the present invention.

5

Preferably, X represents oxygen.

The compounds of formula (I) and their pharmaceutically acceptable salts, enantiomers and racemates have the advantage that they are inhibitors of the enzyme IKK2.

10

The invention further provides a process for the preparation of compounds of formula (I) or a pharmaceutically acceptable salt, enantiomer or racemate thereof.

15

According to the invention there is also provided a compound of formula (I), or a pharmaceutically acceptable salt, enantiomer or racemate thereof, for use as a medicament.

20

Another aspect of the invention provides the use of a compound of formula (I) or a pharmaceutically acceptable salt, enantiomer or racemate thereof, in the manufacture of a medicament, for the treatment or prophylaxis of diseases or conditions in which inhibition of IKK2 activity is beneficial.

25

A more particular aspect of the invention provides the use of a compound of formula (I) or a pharmaceutically acceptable salt, enantiomer or racemate thereof, in the manufacture of a medicament, for the treatment or prophylaxis of inflammatory disease.


30

According to the invention, there is also provided a method of treating, or reducing the risk of, diseases or conditions in which inhibition of IKK2 activity is beneficial which comprises administering to a person suffering from or at risk of, said disease or condition, a therapeutically effective amount of a compound of formula (I), or a pharmaceutically acceptable salt, enantiomer or racemate thereof.

More particularly, there is also provided a method of treating, or reducing the risk of, inflammatory disease in a person suffering from or at risk of, said disease, wherein the method comprises administering to the person a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt, enantiomer or racemate thereof.

In formula (I) the group A is a 5-membered heteroaromatic ring containing one or two heteroatoms selected independently from oxygen, nitrogen or sulfur. Preferably A is substituted as shown below in formula (Ia) where B and D are selected from CR^2 , S, O and

NR²⁵ where R²⁵ is hydrogen or C₁-C₆ alkyl:



Preferred A groups include thiophene, furan, pyrrole, imidazole, thiazole and oxazole. It is particularly preferred that ring A represents thiophene.

15

Suitably the group R¹ is a phenyl group or a 5- to 7-membered heteroaromatic ring containing one to three heteroatoms selected independently from oxygen, nitrogen or sulfur; said phenyl or heteroaromatic ring being optionally substituted by one or more substituents selected from halogen, cyano, nitro, -NR³R⁴, -CONR⁵R⁶, -COOR⁷, -NR⁸COR⁹, -SR¹⁰, -S(O)_mR¹⁰, -SO₂NR⁵R⁶, -NR⁸SO₂R¹⁰, C₁-C₆ alkyl, trifluoromethyl, -(CH₂)_nR¹¹, -O(CH₂)_nR¹¹ or -OR¹². Preferred substituents are halogen, cyano, -NR³R⁴, -SO₂R¹⁰, trifluoromethyl, -O(CH₂)_nR¹¹ or -OR¹². In one preferred embodiment, R¹ represents optionally substituted phenyl. In another preferred embodiment, R¹ represents an optionally substituted 5- or 6-membered heteroaromatic ring containing one or two heteroatoms selected independently from oxygen, nitrogen or sulfur.

25

When R^{11} is $NR^{21}R^{22}$ and R^{21} and R^{22} together with the nitrogen atom to which they are attached form a 5- or 6-membered saturated ring, preferred examples of such rings include morpholine, pyrrolidine and piperidine rings. When R^{11} is $NR^{21}R^{22}$ and R^{21} and R^{22} are alkyl, these alkyl groups are preferably methyl.

Particularly advantageous compounds of formula (I) are those in which R^1 represents optionally substituted phenyl. More preferably R^1 represents phenyl or phenyl substituted by halogen, methoxy, hydroxy, $OCH_2CH_2NMe_2$, $OCH_2CH_2CH_2NMe_2$, morphinolyethoxy, pyrrolidinylethoxy and piperidylethoxy.

It is preferred that the group R^2 in formula (I) represents H, halogen or C_1 - C_2 -alkyl. It is more preferred that the group R^2 represents H or methyl. It is even more preferred that the group R^2 in formula (I) represents H.

Particularly preferred compounds of the invention include those exemplified herein:

- 3-[(aminocarbonyl)amino]-5-phenyl-2-thiophenecarboxamide;
- 3-[(aminocarbonyl)amino]-5-(3-chlorophenyl)-2-thiophenecarboxamide;
- 3-[(aminocarbonyl)amino]-5-(4-fluorophenyl)-2-thiophenecarboxamide;
- 3-[(aminocarbonyl)amino]-5-(4-chlorophenyl)-2-thiophenecarboxamide;
- 3-[(aminocarbonyl)amino]-5-(4-isobutylphenyl)-2-thiophenecarboxamide;
- 3-[(aminocarbonyl)amino]-5-(2-thienyl)-2-thiophenecarboxamide;
- 3-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-2-thiophenecarboxamide;
- 3-[(aminocarbonyl)amino]-5-(3-thienyl)-2-thiophenecarboxamide;
- 3-[(aminocarbonyl)amino]-5-(3-hydroxyphenyl)-2-thiophenecarboxamide;
- 3-[(aminocarbonyl)amino]-5-(2-chlorophenyl)-2-thiophenecarboxamide;

- 3-[(aminocarbonyl)amino]-5-(2-methoxyphenyl)-2-thiophenecarboxamide;
- 3-[(aminocarbonyl)amino]-5-{2-[2-(dimethylamino)ethoxy]phenyl}-2-thiophenecarboxamide;
- 3-[(aminocarbonyl)amino]-5-{4-[2-(dimethylamino)ethoxy]phenyl}-2-thiophenecarboxamide;
- 5 3-[(aminocarbonyl)amino]-5-(3-methoxyphenyl)-2-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-5-phenyl-3-thiophenecarboxamide;
- 3-[(aminocarbonyl)amino]-5-{4-[2-(1-morpholinyl)ethoxy]phenyl}-2-thiophenecarboxamide;
- 10 3-[(aminocarbonyl)amino]-5-{4-[2-(1-pyrrolidinyl)ethoxy]phenyl}-2-thiophenecarboxamide;
- 3-[(aminocarbonyl)amino]-5-{4-[2-(1-piperidinyl)ethoxy]phenyl}-2-thiophenecarboxamide;
- 3-[(aminocarbonyl)amino]-5-{4-[3-(dimethylamino)propoxy]phenyl}-2-thiophenecarboxamide;
- 15 3-[(aminocarbonyl)amino]-5-{3-[2-(dimethylamino)ethoxy]phenyl}-2-thiophenecarboxamide;
- 3-[(aminocarbonyl)amino]-5-{3-[2-(1-morpholinyl)ethoxy]phenyl}-2-thiophenecarboxamide;
- 20 3-[(aminocarbonyl)amino]-5-{3-[2-(1-pyrrolidinyl)ethoxy]phenyl}-2-thiophenecarboxamide;
- 3-[(aminocarbonyl)amino]-5-{3-[2-(1-piperidinyl)ethoxy]phenyl}-2-thiophenecarboxamide;

- 3-[(aminocarbonyl)amino]-5-{3-[3-(dimethylamino)propoxy]phenyl}-2-thiophenecarboxamide;
- 3-[(aminocarbonyl)amino]-5-{2-[2-(1-morpholinyl)ethoxy]phenyl}-2-thiophenecarboxamide;
- 5 3-[(aminocarbonyl)amino]-5-{2-[2-(1-pyrrolidinyl)ethoxy]phenyl}-2-thiophenecarboxamide;
- 3-[(aminocarbonyl)amino]-5-{2-[2-(1-piperidinyl)ethoxy]phenyl}-2-thiophenecarboxamide;
- 3-[(aminocarbonyl)amino]-5-{2-[3-(dimethylamino)propoxy]phenyl}-2-
- 10 thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-4-methyl-5-(4-chlorophenyl)-3-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-4-methyl-5-(4-methylphenyl)-3-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-4-ethyl-5-phenyl-3-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-4-methyl-5-(4-methoxyphenyl)-3-thiophenecarboxamide;
- 15 2-[(aminocarbonyl)amino]-4-methyl-5-(4-fluorophenyl)-3-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-4-methyl-5-(3-fluorophenyl)-3-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-4-methyl-5-(3-methoxyphenyl)-3-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-4-methyl-5-(3-chloro-4-methoxyphenyl)-3-
- thiophenecarboxamide;
- 20 2-[(aminocarbonyl)amino]-4-methyl-5-(2-chlorophenyl)-3-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-4-methyl-5-(3-trifluoromethylphenyl)-3-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-4-methyl-5-(3-methyl-4-methoxyphenyl)-3-
- thiophenecarboxamide;

- 2-[(aminocarbonyl)amino]-4-methyl-5-(3,5-dimethoxyphenyl)-3-thiophenecarboxamide;
2-[(aminocarbonyl)amino]-4-methyl-5-(2,3-dimethoxyphenyl)-3-thiophenecarboxamide;
2-[(aminocarbonyl)amino]-4-methyl-5-(4-isopropylphenyl)-3-thiophenecarboxamide;
2-[(aminocarbonyl)amino]-4-methyl-5-(3,4,5-trimethoxyphenyl)-3-thiophenecarboxamide;
5 2-[(aminocarbonyl)amino]-4-methyl-5-(2-pyridyl)-3-thiophenecarboxamide;
2-[(aminocarbonyl)amino]-5-[2-(5-methoxypyridyl)]-4-methyl-3-thiophenecarboxamide;
2-[(aminocarbonyl)amino]-4-methyl-5-(4-pyrimidyl)-3-thiophenecarboxamide;
2-[(aminocarbonyl)amino]-4-methyl-5-(2-pyrazinyl)-3-thiophenecarboxamide;
2-[(aminocarbonyl)amino]-4-methyl-5-(3,4-dichlorophenyl)-3-thiophenecarboxamide;
10 2-[(aminocarbonyl)amino]-4-methyl-5-(4-cyanophenyl)-3-thiophenecarboxamide;
2-[(aminocarbonyl)amino]-4-methyl-5-(4-hydroxyphenyl)-3-thiophenecarboxamide;
2-[(aminocarbonyl)amino]-4-methyl-5-(4-[2-(1-piperidinyl)ethoxy]phenyl)-3-
thiophenecarboxamide;
2-[(aminocarbonyl)amino]-4-methyl-5-(4-[2-(diethylamino)ethoxy]phenyl)-3-
15 thiophenecarboxamide;
2-[(aminocarbonyl)amino]-4-methyl-5-(2-furyl)-3-thiophenecarboxamide;
2-[(aminocarbonyl)amino]-4-trifluoromethyl-5-phenyl-3-thiophenecarboxamide;
2-[(aminocarbonyl)amino]-4-methyl-5-(2-(4-methylthiazolyl))-3-thiophenecarboxamide;
2-[(aminocarbonyl)amino]-4-methyl-5-phenyl-3-thiophenecarboxamide;
20 2-[(aminocarbonyl)amino]-4-methyl-5-(3-methyl-isoxazol-5-yl)-3-thiophenecarboxamide;
2-[(aminocarbonyl)amino]-5-(4-cyanophenyl)-3-thiophenecarboxamide;
2-[(aminocarbonyl)amino]-5-(4-trifluoromethylphenyl)-3-thiophenecarboxamide;
2-[(aminocarbonyl)amino]-5-(2,4-difluorophenyl)-3-thiophenecarboxamide;

- 2-[(aminocarbonyl)amino]-5-(2-pyridyl)-3-thiophenecarboxamide;
2-[(aminocarbonyl)amino]-5-(3-pyridyl)-3-thiophenecarboxamide;
2-[(aminocarbonyl)amino]-5-[5-(2-methoxypyridyl)]-3-thiophenecarboxamide;
2-[(aminocarbonyl)amino]-5-[5-(2,4-dimethoxypyrimidyl)]-3-thiophenecarboxamide;
5 2-[(aminocarbonyl)amino]-5-(4-hydroxyphenyl)-3-thiophenecarboxamide;
2-[(aminocarbonyl)amino]-5-(4-chlorophenyl)-3-thiophenecarboxamide;
2-[(aminocarbonyl)amino]-5-(4-methanesulphonylphenyl)-3-thiophenecarboxamide;
2-[(aminocarbonyl)amino]-5-(2-(N-t-butoxycarbonyl)pyrrolyl)-3-thiophenecarboxamide;
2-[(aminocarbonyl)amino]-5-(2-(5-cyanothienyl))-3-thiophenecarboxamide;
10 2-[(aminocarbonyl)amino]-5-(3,5-dimethyl-isoxazol-4-yl)-3-thiophenecarboxamide;
2-[(aminocarbonyl)amino]-5-(3-furyl)-3-thiophenecarboxamide;
2-[(aminocarbonyl)amino]-5-(2-pyrrolyl)-3-thiophenecarboxamide;
2-[(aminocarbonyl)amino]-5-(5-pyrimidinyl)-3-thiophenecarboxamide;
2-[(aminocarbonyl)amino]-5-(2-(5-chlorothienyl))-3-thiophenecarboxamide;
15 2-[(aminocarbonyl)amino]-5-[2-(5-trifluoromethylpyridyl)]-3-thiophenecarboxamide;
2-[(aminocarbonyl)amino]-5-[2-(5-bromopyridyl)]-3-thiophenecarboxamide;
2-[(aminocarbonyl)amino]-5-(2-(5-cyanofuryl))-3-thiophenecarboxamide;
2-[(aminocarbonyl)amino]-5-(4-[2-(1-piperidinyl)ethoxy]phenyl)-3-
thiophenecarboxamide;
20 2-[(aminocarbonyl)amino]-5-(4-[2-(1-(2,2,6,6-tetramethyl)piperidinyl)ethoxy]phenyl)-3-
thiophenecarboxamide;
2-[(aminocarbonyl)amino]-5-(4-(thiazol-4-yl-methoxy)phenyl)-3-thiophenecarboxamide;

2-[(aminocarbonyl)amino]-5-(4-[2-(dimethylamino)ethoxy]phenyl)-3-thiophenecarboxamide;

2-[(aminocarbonyl)amino]-5-(4-[2-(diethylamino)ethoxy]phenyl)-3-thiophenecarboxamide;

5 2-[(aminocarbonyl)amino]-5-(4-[2-(1-morpholinyl)ethoxy]phenyl)-3-thiophenecarboxamide;

2-[(aminocarbonyl)amino]-5-(2-furyl)-3-thiophenecarboxamide;

2-[(aminocarbonyl)amino]-5-(2-(5-methylfuryl))-3-thiophenecarboxamide;

5-[(aminocarbonyl)amino]-2-(3,5-dichlorophenyl)-1,3-oxazole-4-carboxamide;

10 5-[(aminocarbonyl)amino]-2-(4-trifluoromethylphenyl)-1,3-oxazole-4-carboxamide;

2-[(aminothiocabonyl)amino]-5-phenyl-3-thiophenecarboxamide;

and pharmaceutically acceptable salts and solvates thereof.

Unless otherwise indicated, the term " C_1 - C_6 alkyl" referred to herein denotes a straight or
15 branched chain alkyl group having from 1 to 6 carbon atoms. Examples of such groups include methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl and t-butyl. The term " C_1 - C_2 alkyl" is to be interpreted analogously.

Unless otherwise indicated, the term " C_2 - C_3 alkenyl" referred to herein denotes a straight or
20 branched chain alkyl group having 2 or 3 carbon atoms incorporating at least one carbon-carbon double bond. Examples of such groups include ethenyl and propenyl.

Unless otherwise indicated, the term " C_2 - C_3 alkynyl" referred to herein denotes a straight
chain alkyl group having 2 or 3 carbon atoms incorporating one carbon-carbon triple bond.
25 Examples of such groups include ethynyl and propynyl.

Unless otherwise indicated, the term "C₁-C₄ alkoxy" referred to herein denotes a straight or branched chain alkoxy group having 1 to 4 carbon atoms. Examples of such groups include methoxy, ethoxy and isopropoxy. The term "C₁-C₂ alkoxy" is to be interpreted analogously.

- 5 Unless otherwise indicated, the term "C₁-C₂ alkanoyl" referred to herein denotes a formyl or acetyl group.

Unless otherwise indicated, the term "halogen" referred to herein denotes fluoro, chloro, bromo and iodo.


10

Examples of a 5- to 7-membered heteroaromatic ring containing one to three heteroatoms selected independently from oxygen, nitrogen or sulfur include furan, thiophene, pyrrole, oxazole, isoxazole, thiazole, isothiazole, imidazole, pyrazole, triazole, pyridine, pyridazine, pyrimidine and pyrazine.

15

According to the invention there is also provided a process for the preparation of a compound of formula (I) or a pharmaceutically acceptable salt, enantiomer or racemate thereof which comprises:

- 20 (a) reaction of a compound of formula (II):




wherein A, R¹ and R² are as defined in formula (I) with an isocyanate (X = O) or an isothiocyanate (X = S); or

25

- (b) reaction of compound of formula (III) with a compound of formula (IV)

R¹-Metal

(III)



(IV)


wherein A, X, R¹ and R² are as defined in formula (I) and LG represents a leaving group; or

(c) reaction of compound of formula (V) with a compound of formula (VI)

5

R¹-LG

(V)



(VI)

wherein A, X, R¹ and R² are as defined in formula (I) and LG represents a leaving group;

and where necessary converting the resultant compound of formula (I), or another salt thereof, into a pharmaceutically acceptable salt thereof; or converting the resultant compound of formula (I) into a further compound of formula (I); and where desired converting the resultant compound of formula (I) into an optical isomer thereof.

In process (a), suitable isocyanate reagents include trimethylsilylisocyanate, trimethylsilylisothiocyanate, chlorosulphonylisocyanate, trichloroacetylisocyanate and sodium isocyanate. The reaction with trimethylsilylisocyanate or trimethylsilylisothiocyanate can be carried out in a solvent such as dichloromethane/dimethylformamide at a suitable elevated temperature, for example, at the reflux temperature of the reaction mixture. The reaction with chlorosulphonylisocyanate can be carried out in a solvent such as toluene at ambient temperature. The reaction with

sodium isocyanate can be carried out in a suitable solvent system such as aqueous acetic acid at ambient temperature. The trichloroacetylisocyanate reaction can be carried out in a suitable solvent system such as acetonitrile at ambient temperature, and subsequently treating the mixture with ammonia to give compounds of the general formula (I).

5

In processes (b) and (c), the compounds of formulae (III) and (IV) or of formulae (V) and (VI) are reacted together under catalysis provided by a complex of a transition metal such as palladium or nickel. In compounds of formulae (III) and (VI), under appropriate conditions, "metal" can be a metal or semi-metal such as magnesium, zinc, copper, tin,
10 silicon, zirconium, aluminium or boron. Suitable leaving groups include iodo, bromo, chloro, triflate or phosphonate.


It will be appreciated by those skilled in the art that in the processes of the present invention certain functional groups such as hydroxyl or amino groups in the starting
15 reagents or intermediate compounds may need to be protected by protecting groups. Thus, the preparation of the compounds of formula (I) may involve, at an appropriate stage, the addition and removal of one or more protecting groups.

The protection and deprotection of functional groups is fully described in 'Protective
20 Groups in Organic Chemistry', edited by J. W. F. McOmie, Plenum Press (1973), and 'Protective Groups in Organic Synthesis', 2nd edition, T. W. Greene & P. G. M. Wuts, Wiley-Interscience (1991).

The present invention includes compounds of formula (I) in the form of salts, in particular
25 acid addition salts. Suitable salts include those formed with both organic and inorganic acids. Such acid addition salts will normally be pharmaceutically acceptable although salts of non-pharmaceutically acceptable acids may be of utility in the preparation and purification of the compound in question. Thus, preferred salts include those formed from hydrochloric, hydrobromic, sulphuric, phosphoric, citric, tartaric, lactic, pyruvic, acetic,
30 succinic, fumaric, maleic, methanesulphonic and benzenesulphonic acids.

Salts of compounds of formula (I) may be formed by reacting the free base, or a salt, enantiomer or racemate thereof, with one or more equivalents of the appropriate acid. The reaction may be carried out in a solvent or medium in which the salt is insoluble or in a solvent in which the salt is soluble, for example, water, dioxane, ethanol, tetrahydrofuran or diethyl ether, or a mixture of solvents, which may be removed *in vacuo* or by freeze drying. The reaction may also be a metathetical process or it may be carried out on an ion exchange resin.


Compounds of formula (II) can be prepared by standard chemistry described in the literature [for example, J. Het. Chem. 36, 333 (1999)] or by reaction of compounds of formula (VII):



(VII)

where A, R^1 and R^2 are as defined in formula (I), and L represents a leaving group, with ammonia. Suitable groups L include halogen, in particular chloro.

Compounds of formula (VII) where L is halo can be prepared from the corresponding compound of formula (VIII):



(VIII)

where A, R^1 and R^2 are as defined in formula (I), by treating with a halogenating agent such as thionyl chloride.

Compounds of formulae (III), (IV), (V), (VI) and (VIII) are commercially available or can be prepared using standard chemistry as exemplified herein.

5 Certain novel intermediate compounds form a further aspect of the invention.

The compounds of formula (I) have activity as pharmaceuticals, in particular as IKK2 enzyme inhibitors, and may be used in the treatment (therapeutic or prophylactic) of conditions/diseases in human and non-human animals in which inhibition of IKK2 is
10 beneficial. Examples of such conditions/diseases include inflammatory diseases or diseases with an inflammatory component. Particular diseases include inflammatory arthritides including rheumatoid arthritis, osteoarthritis, spondylitis, Reiters syndrome, psoriatic arthritis, lupus and bone resorptive disease; multiple sclerosis, inflammatory bowel disease including Crohn's disease; asthma, chronic obstructive pulmonary disease,
15 emphysema, rhinitis, myasthenia gravis, Graves' disease, allograft rejection, psoriasis, dermatitis, allergic disorders, immune complex diseases, cachexia, ARDS, toxic shock, cardiovascular disorders, heart failure, myocardial infarcts, atherosclerosis, reperfusion injury, AIDS and cancer.

20 Thus, the present invention provides a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as hereinbefore defined for use in therapy.

In a further aspect, the present invention provides the use of a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as hereinbefore defined in the
25 manufacture of a medicament for use in therapy.

In a still further aspect, the present invention provides the use of a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as hereinbefore defined in the manufacture of a medicament for the treatment of diseases or conditions in which
30 modulation of the IKK2 enzyme activity is beneficial.

In the context of the present specification, the term "therapy" also includes "prophylaxis" unless there are specific indications to the contrary. The terms "therapeutic" and "therapeutically" should be construed accordingly.

5 Prophylaxis is expected to be particularly relevant to the treatment of persons who have suffered a previous episode of, or are otherwise considered to be at increased risk of, the disease or condition in question. Persons at risk of developing a particular disease or condition generally include those having a family history of the disease or condition, or those who have been identified by genetic testing or screening to be particularly
10 susceptible to developing the disease or condition.

The invention still further provides a method of treating an IKK2 mediated disease which comprises administering to a patient a therapeutically effective amount of a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as hereinbefore
15 defined.

The invention also provides a method of treating an inflammatory disease, especially asthma, rheumatoid arthritis or multiple sclerosis, in a patient suffering from, or at risk of, said disease, which comprises administering to the patient a therapeutically effective
20 amount of a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as hereinbefore defined.

For the above-mentioned therapeutic uses the dosage administered will, of course, vary with the compound employed, the mode of administration, the treatment desired and the
25 disorder indicated.

The compounds of formula (I) and pharmaceutically acceptable salts and solvates thereof may be used on their own but will generally be administered in the form of a pharmaceutical composition in which the formula (I) compound/salt/solvate (active
30 ingredient) is in association with a pharmaceutically acceptable adjuvant, diluent or carrier. Depending on the mode of administration, the pharmaceutical composition will preferably

comprise from 0.05 to 99 %w (per cent by weight), more preferably from 0.05 to 80 %w, still more preferably from 0.10 to 70 %w, and even more preferably from 0.10 to 50 %w, of active ingredient, all percentages by weight being based on total composition.

- 5 The present invention also provides a pharmaceutical composition comprising a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as hereinbefore defined, in association with a pharmaceutically acceptable adjuvant, diluent or carrier.

The invention further provides a process for the preparation of a pharmaceutical
10 composition of the invention which comprises mixing a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as hereinbefore defined, with a pharmaceutically acceptable adjuvant, diluent or carrier.

The pharmaceutical compositions may be administered topically (e.g. to the lung and/or
15 airways or to the skin) in the form of solutions, suspensions, heptafluoroalkane aerosols and dry powder formulations; or systemically, e.g. by oral administration in the form of tablets, capsules, syrups, powders or granules, or by parenteral administration in the form of solutions or suspensions, or by subcutaneous administration or by rectal administration in the form of suppositories or transdermally. Conventional procedures for the selection
20 and preparation of suitable pharmaceutical formulations are described in, for example, "Pharmaceuticals - The Science of Dosage Form Designs", M. E. Aulton, Churchill Livingstone, 1988.

The invention is illustrated, but in no way limited, by the following examples:

25

EXAMPLE 1

3-[(Aminocarbonyl)amino]-5-phenyl-2-thiophenecarboxamide

3-Amino-5-phenyl-2-thiophenecarboxamide (0.5 g), trimethylsilylisocyanate (3 mL),
30 dichloromethane (15 mL) and dimethylformamide (3 mL) were heated at reflux for 3 days.

The reaction mixture was cooled and the resulting solid was filtered off, washed with methanol and then ether to give the title urea (0.39 g).

m.p. >300 °C.

¹H NMR (DMSO-D₆) 10.06 (1H, s), 8.25 (1H, s), 7.62 (2H, d), 7.50-7.37 (5H, m), 6.63
5 (2H, s).

EXAMPLE 2

3-[(Aminocarbonyl)amino]-5-(3-chlorophenyl)-2-thiophenecarboxamide

10 a) Methyl 3-amino-5-(3-chlorophenyl)-2-thiophenecarboxylate

Phosphorous oxychloride (6.7 mL) was added to dimethylformamide (11 mL) with ice cooling to keep the internal temperature below 25 °C. After 20 minutes, (3-chlorophenyl)ethanone (5 g) was added portionwise keeping the internal temperature below 30 °C. The reaction mixture was heated to 50 °C and then treated cautiously with
15 hydroxylamine hydrochloride (10 g). The reaction mixture was stirred for 20 minutes at room temperature and water (50 mL) was added. After a further 30 minutes, the reaction mixture was extracted three times with ethyl acetate. The combined extracts were washed with brine, dried (MgSO₄) and evaporated to give an oil. This oil was dissolved in methanol (50 mL) and treated with methyl mercaptoacetate (2.7 mL) and sodium
20 methoxide (7.3 mL of a 25% solution in methanol). After reflux for 1 h, the cooled reaction mixture was reduced to one third volume and water was added. The reaction mixture was extracted three times with ethyl acetate. The combined extracts were dried (MgSO₄), the solvent was evaporated and the residue was chromatographed on silica eluting with dichloromethane/isohexane mixtures to give the sub-title ester (2.0 g).

25 m.p. 105-6 °C.

MS (EI) 267 (M)⁺.

¹H NMR (DMSO-D₆) 7.68 (1H, s); 7.60 (1H, m); 7.48 (2H, m); 7.02 (1H, s); 6.60 (2H, s);
3.74 (3H, s).

30 b) 3-Amino-5-(3-chlorophenyl)-2-thiophenecarboxylic acid

Methyl 3-amino-5-(3-chlorophenyl)-2-thiophenecarboxylate (1.0 g), 2M sodium hydroxide (2 mL) and methanol (10 mL) were heated at 70 °C for 2 days. The methanol was evaporated off and the residue was acidified with 2M hydrochloric acid. Extraction into ethyl acetate followed by drying (MgSO₄) and evaporation of the solvent gave the sub-title acid (0.8 g).

MS (APCI) 252 (M+H)⁺.

¹H NMR (DMSO-D₆) 7.62 (1H, d); 7.60 (1H, m); 7.43 (2H, m); 7.02 (1H, s).

c) 3-Amino-5-(3-chlorophenyl)-2-thiophenecarboxamide

3-Amino-5-(3-chlorophenyl)-2-thiophenecarboxylic acid (0.8 g) and thionyl chloride (20 mL) were heated at reflux for 1 h. After cooling, the excess thionyl chloride was evaporated off and final traces were removed by azeotropeing with toluene. The residue was dissolved in acetonitrile (50 mL) and ammonia (d 0.88, 10 mL) was added. After stirring for 1 h, the solvent was evaporated and the residue chromatographed on silica eluting with ethyl acetate/dichloromethane mixtures. Trituration with ether gave the sub-title amide (0.48 g).

m.p. 164-5 °C

MS (APCI) 253 (M+H)⁺.

¹H NMR (DMSO-D₆) 7.62 (1H, d); 7.55 (1H, dd); 7.45 (2H, m); 7.02 (1H, s); 6.98 (2H, s); 6.50 (2H, s).

d) 3-[(Aminocarbonyl)amino]-5-(3-chlorophenyl)-2-thiophenecarboxamide

Prepared by the method of Example 1 from 3-amino-5-(3-chlorophenyl)-2-thiophenecarboxamide and trimethylsilylisocyanate.

m.p. >300 °C.

MS (APCI) 253 (M+H)⁺.

¹H NMR (DMSO-D₆) 10.03 (1H, s); 8.30 (1H, s); 7.62 (1H, d); 7.60-7.40 (4H, m); 7.30-7.00 (1H, m); 6.70 (2H, s).

EXAMPLE 33-[(Aminocarbonyl)amino]-5-(4-fluorophenyl)-2-thiophenecarboxamide

- Sodium isocyanate (1.08 g) was added portionwise to a stirred suspension of 3-amino-5-(4-fluorophenyl)-2-thiophenecarboxamide (3.2 g) in acetic acid (150 mL) and water (90 mL). After 20 h, the solid was filtered off and washed with water, methanol and ether. Recrystallisation from methanol/dimethylsulphoxide gave the title urea (0.5 g) as a 1:1 dimethylsulphoxide solvate.
- m.p. >320 °C.
- MS (APCI) 278 (M-H)⁺.
- ¹H NMR (DMSO-D₆) 10.07 (1H, s); 8.22 (1H, s); 7.67 (2H, t); 7.40 (2H, s); 7.29 (2H, t); 6.65 (2H, s).

EXAMPLE 3a

- 15 3-[(Aminocarbonyl)amino]-5-(4-chlorophenyl)-2-thiophenecarboxamide

- Prepared by the method of Example 3 from 3-amino-5-(4-chlorophenyl)-2-thiophenecarboxamide.
- 20 MS (ES) 296 (M+H)⁺.
- ¹H NMR (DMSO-D₆) 11.03 (1H, s), 8.2 (1H, s), 7.6 (2H, d), 7.5 (2H, d), 7.4 (2H, s), 6.8 (2H, s).

EXAMPLE 3b

- 25 3-[(Aminocarbonyl)amino]-5-(4-isobutylphenyl)-2-thiophenecarboxamide

Prepared by the method of Example 3 from 3-amino-5-(4-isobutylphenyl)-2-thiophenecarboxamide.

MS (ES) 318 (M+H)⁺.

¹H NMR (DMSO-D₆) 11.03 (1H, s), 8.2 (1H, s), 7.5 (2H, m), 7.4 (2H, s), 7.2 (2H, m), 6.6 (2H, s), 2.4 (1H, m), 1.8 (2H, m), 0.8 (6H, m).

EXAMPLE 3c

5 3-[(Aminocarbonyl)amino]-5-(2-thienyl)-2-thiophenecarboxamide

Prepared by the method of Example 3 from 3-amino-5-(2-thienyl)-2-thiophenecarboxamide.

MS (ES) 266 (M-H).

10 ¹H NMR (DMSO-D₆) 10.03 (1H, s), 8.05 (1H, s), 7.6 (1H, d), 7.4 (3H, m), 7.1 (1H, t), 6.6 (2H, s).

EXAMPLE 4

15 3-[(Aminocarbonyl)amino]-5-(4-methoxyphenyl)-2-thiophenecarboxamide

Prepared by the method of Example 1 from 3-amino-5-(4-methoxyphenyl)-2-thiophenecarboxamide and trimethylsilylisocyanate.

m.p. >300 °C.

MS (APCI) 292 (M+H)⁺.

20 ¹H NMR (DMSO-D₆) 10.06 (1H, s); 8.12 (1H, s); 7.55 (2H, d); 7.37 (2H, s); 7.03 (2H, d); 6.61 (2H, s); 3.80 (3H, s).

EXAMPLE 5

25 3-[(Aminocarbonyl)amino]-5-(3-thienyl)-2-thiophenecarboxamide

Prepared by the method of Example 1 from 3-amino-5-(3-thienyl)-2-thiophenecarboxamide and trimethylsilylisocyanate.

¹H NMR (DMSO-D₆) 10.0 (1H, s), 8.05 (1H, s), 7.8 (1H, d), 7.65 (1H, m), 7.4 (3H, m), 6.6 (2H, s).

EXAMPLE 63-[(Aminocarbonyl)amino]-5-(3-hydroxyphenyl)-2-thiophenecarboxamide

5 3-Amino-5-(3-methoxyphenyl)-2-thiophenecarboxamide (0.5 g), trimethylsilylisocyanate (2 mL), dimethylformamide (2 mL) and dichloromethane were heated at reflux for 3 days. After cooling the solid was filtered off, suspended in dichloromethane (100 mL) and treated with boron tribromide (5 mL of a 1M solution in dichloromethane). After 3 days, methanol (50 mL) was added. After 1 h, the solvent was evaporated and the residue was
10 triturated with 2M hydrochloric acid. The title urea was filtered off (0.35 g).
m.p. >300 °C.

MS (APCI) 278 (M+H)⁺.

¹H NMR (DMSO-D₆) 10.05 (1H, s); 9.71 (1H, s); 8.19 (1H, s); 7.41 (2H, m); 7.26 (1H, t);
7.03 (2H, m); 6.79 (1H, dd); 6.62 (2H, s).

15

EXAMPLE 73-[(Aminocarbonyl)amino]-5-(2-chlorophenyl)-2-thiophenecarboxamidea) 3-Amino-5-(2-chlorophenyl)-2-thiophenecarboxylic acid

20 Prepared by the method of Example 2(b) from methyl 3-amino-5-(2-chlorophenyl)-2-thiophenecarboxylate.

MS (APCI) 252 (M+H)⁺.

¹H NMR (DMSO-D₆) 7.60 (2H, m); 7.40 (2H, m); 6.92 (1H, s).

25 b) 3-Amino-5-(2-chlorophenyl)-2-thiophenecarboxamide

Prepared by the method of Example 2(c) from 3-amino-5-(2-chlorophenyl)-2-thiophenecarboxylic acid.

m.p. 87-89 °C.

MS (APCI) 253 (M+H)⁺.

30 ¹H NMR (DMSO-D₆) 7.60 (2H, m); 7.40 (2H, m); 7.00 (2H, s); 6.90 (1H, s); 6.42 (2H, s).

c) 3-[(Aminocarbonyl)amino]-5-(2-chlorophenyl)-2-thiophenecarboxamide

Prepared by the method of Example 1 from 3-amino-5-(2-chlorophenyl)-2-thiophenecarboxamide and trimethylsilylisocyanate.

5 m.p. >300 °C.

MS (APCI) 296 (M+H)⁺.

¹H NMR (DMSO-D₆) 7.34 (2H, s); 6.80 (2H, m); 6.70 (2H, m); 6.52 (4H, m).

EXAMPLE 8

10 3-[(Aminocarbonyl)amino]-5-(2-methoxyphenyl)-2-thiophenecarboxamide

a) Methyl 3-amino-5-(2-methoxyphenyl)-2-thiophenecarboxylate

Prepared by the method of Example 2(a) from (2-methoxyphenyl)ethanone.

m.p. 119-20 °C.

15 MS (APCI) 264 (M+H)⁺.

¹H NMR (DMSO-D₆) 7.62 (1H, dd); 7.40 (1H, t); 7.18 (1H, d); 7.05 (1H, s); 7.02 (1H, t); 6.45 (2H, s); 3.95 (3H, s); 3.75 (3H, s).

b) 3-Amino-5-(2-methoxyphenyl)-2-thiophenecarboxylic acid

20 Prepared by the method of Example 2(b) from methyl 3-amino-5-(2-methoxyphenyl)-2-thiophenecarboxylate and used directly for step (c).

c) 3-Amino-5-(2-methoxyphenyl)-2-thiophenecarboxamide

Prepared by the method of Example 2(c) from 3-amino-5-(2-methoxyphenyl)-2-thiophenecarboxylic acid and used directly for step (d).

25

d) 3-[(Aminocarbonyl)amino]-5-(2-methoxyphenyl)-2-thiophenecarboxamide

Prepared by the method of Example 1 from 3-amino-5-(2-methoxyphenyl)-2-thiophenecarboxamide and trimethylsilylisocyanate.

30 m.p. >300 °C.

¹H NMR (DMSO-D₆) 10.01 (1H, s); 8.33 (1H, s); 7.62 (1H, dd); 7.40-7.00 (5H, m); 6.57 (2H, s); 3.90 (3H, s).

EXAMPLE 9

5 3-[(Aminocarbonyl)amino]-5-{2-[2-(dimethylamino)ethoxy]phenyl}-2-thiophenecarboxamide

a) 3-[(Aminocarbonyl)amino]-5-(2-hydroxyphenyl)-2-thiophenecarboxamide

3-[(Aminocarbonyl)amino]-5-(2-methoxyphenyl)-2-thiophenecarboxamide (0.1 g), boron
10 tribromide (2 ml of a 1M solution in dichloromethane) and dichloromethane (10 mL) were stirred at room temperature for 16 h. Methanol (5 mL) was added and after 1 h, the solvent was evaporated. 2M Hydrochloric acid (10 mL) was added and, after stirring for 1 h, the phenol was filtered off and used directly in step (b).

15 b) 3-[(Aminocarbonyl)amino]-5-{2-[2-(dimethylamino)ethoxy]phenyl}-2-thiophenecarboxamide

The phenol (0.05 g), potassium carbonate (0.05 g) and (2-chloroethyl)dimethylamine hydrochloride (0.03 g) in dimethylformamide (2 mL) were stirred at 80 °C for 24 h. The cooled reaction was poured onto ethyl acetate and brine. The aqueous layer was separated
20 and washed twice with ethyl acetate. The combined organic extracts were washed with brine, dried (MgSO₄) and the solvent was evaporated. Chromatography on silica eluting with dichloromethane/methanol mixtures gave the title compound (6 mg).
m.p. 180 °C.

MS (APCI) 349 (M+H)⁺.

25 ¹H NMR (DMSO-D₆) 10.00 (1H, s); 8.40 (1H, s); 7.62 (1H, dd); 7.38 (3H, m); 7.20 (1H, d); 7.05 (1H, t); 6.60 (2H, s); 4.20 (2H, t); 2.80 (2H, t); 2.50 (6H, s).

EXAMPLE 10

30 3-[(Aminocarbonyl)amino]-5-{4-[2-(dimethylamino)ethoxy]phenyl}-2-thiophenecarboxamide

a) 3-[(Aminocarbonyl)amino]-5-(4-hydroxyphenyl)-2-thiophenecarboxamide

Prepared by the method of Example 6 from 3-amino-5-(4-methoxyphenyl)-2-thiophenecarboxamide and used directly in step (b).

5 b) 3-[(Aminocarbonyl)amino]-5-{4-[2-(dimethylamino)ethoxy]phenyl}-2-thiophenecarboxamide

Prepared by the method of Example 9(b) from 3-[(aminocarbonyl)amino]-5-(4-hydroxyphenyl)-2-thiophenecarboxamide.

m.p. >300 °C.

10 MS (APCI) 349 (M+H)⁺.

¹H NMR (DMSO-D₆) 10.06 (1H, s); 8.12 (1H, s); 7.53 (2H, d); 7.40 (2H, s); 7.00 (2H, d); 6.60 (2H, s); 4.10 (2H, t); 2.60 (2H, t); 2.20 (6H, s).

EXAMPLE 11

15 3-[(Aminocarbonyl)amino]-5-(3-methoxyphenyl)-2-thiophenecarboxamide

a) Methyl 3-amino-5-(3-methoxyphenyl)-2-thiophenecarboxylate

Prepared by the method of Example 2(a) from (3-methoxyphenyl)ethanone.

m.p. 81-2 °C.

20 MS (APCI) 264 (M+H)⁺.

¹H NMR (DMSO-D₆) 7.40 (1H, t); 7.20 (1H, d); 7.15 (1H, m); 7.00 (2H, m); 6.60 (2H, s); 3.80 (3H, s); 3.70 (3H, s).

b) 3-Amino-5-(3-methoxyphenyl)-2-thiophenecarboxylic acid

25 Prepared by the method of Example 2(b) from methyl 3-amino-5-(3-methoxyphenyl)-2-thiophenecarboxylate and used directly in step (c).

c) 3-Amino-5-(3-methoxyphenyl)-2-thiophenecarboxamide

Prepared by the method of Example 2(c) from 3-amino-5-(3-methoxyphenyl)-2-thiophenecarboxylic acid.

30 m.p. 101-3 °C.

MS (APCI) 249 (M+H)⁺.

¹H NMR (DMSO-D₆) 7.35 (1H, t); 7.20 (1H, d); 7.10 (1H, m); 7.00-6.90 (4H, m); 6.42 (2H, s); 3.80 (3H, s).

d) 3-[(Aminocarbonyl)amino]-5-(3-methoxyphenyl)-2-thiophenecarboxamide

5 Prepared by the method of Example 1 from 3-amino-5-(3-methoxyphenyl)-2-thiophenecarboxamide and trimethylsilylisocyanate.

m.p. 105-6 °C.

MS (EI) 267 (M)⁺.

¹H NMR (DMSO-D₆) 10.05 (1H, s); 8.23 (1H, s), 7.43 (2H, s); 7.39 (1H, t); 7.19 (1H, d);
10 7.10 (1H, s); 6.98 (1H, d); 6.62 (2H, s); 3.82 (3H, s).

EXAMPLE 12

2-[(Aminocarbonyl)amino]-5-phenyl-3-thiophenecarboxamide

15 Chlorosulphonylisocyanate (0.081 mL) was added to a stirred suspension at 0 °C of 2-amino-5-phenyl-3-thiophenecarboxamide (0.2 g) in toluene (10 mL). After stirring for 16 h at room temperature, the solvent was evaporated and the residue dissolved in acetonitrile (20 mL). 10% Sodium bicarbonate solution (2 mL) was added and the mixture

20 extracted three times with ethyl acetate. The combined extracts were dried (MgSO₄) and the solvent was evaporated. Chromatography on silica eluting with methanol/dichloromethane mixtures gave the title urea (0.027 g).

m.p. 395 °C.

MS (APCI) 262 (M+H)⁺.

25 ¹H NMR (DMSO-D₆) 11.01 (1H, s); 7.73 (1H, s); 7.69 (1H, s); 7.58 (1H, s); 7.54 (1H, s); 7.40 (2H, t); 7.35-7.20 (2H, m); 7.00 (2H, s).

The starting 2-amino-5-phenyl-3-thiophenecarboxamide was prepared as follows :

A solution of phenylacetaldehyde (7.2 g), sulphur (1.92 g), cyanoacetamide (5.1 g) and
30 triethylamine (4.53 mL) in dimethylformamide (45 mL) was stirred at room temperature

for 1 h. The resulting solution was added to water (300 mL) and the solid precipitate was filtered off and washed with water. The dried solid was triturated with ether and collected (5.5 g).

5 MS (ES) 219 (M+H)⁺.

¹H NMR (DMSO-D₆) 7.55 (1H, s), 7.4 (2H, m), 7.35 (5H, m), 7.1 (1H, m).

EXAMPLE 13

10 3-[(Aminocarbonyl)amino]-5-{4-[2-(1-morpholinyl)ethoxy]phenyl}-2-thiophenecarboxamide

Prepared by the method of Example 9(b) using the product of Example 10 (a) and N-(2-chloroethyl)morpholine.

MS (EI) 390 (M)⁺.

15

EXAMPLE 14

3-[(Aminocarbonyl)amino]-5-{4-[2-(1-pyrrolidinyl)ethoxy]phenyl}-2-thiophenecarboxamide

20 Prepared by the method of Example 9(b) using the product of Example 10 (a) and N-(2-chloroethyl)pyrrolidine.

MS (EI) 374 (M)⁺.

25

EXAMPLE 15

3-[(Aminocarbonyl)amino]-5-{4-[2-(1-piperidinyl)ethoxy]phenyl}-2-thiophenecarboxamide

30 Prepared by the method of Example 9(b) using the product of Example 10 (a) and N-(2-chloroethyl)piperidine.

MS (EI) 388 (M)⁺.

EXAMPLE 16

3-[(Aminocarbonyl)amino]-5-{4-[3-(dimethylamino)propoxy]phenyl}-2-
thiophenecarboxamide

Prepared by the method of Example 9(b) using the product of Example 10 (a) and N-(3-chloropropyl)dimethylamine.

MS (EI) 362 (M)⁺.

EXAMPLE 17

3-[(Aminocarbonyl)amino]-5-{3-[2-(dimethylamino)ethoxy]phenyl}-2-
thiophenecarboxamide

Prepared by the method of Example 9(b) using the product of Example 6 and N-(2-chloroethyl)dimethylamine.

MS (EI) 348 (M)⁺.

EXAMPLE 18

3-[(Aminocarbonyl)amino]-5-{3-[2-(1-morpholinyl)ethoxy]phenyl}-2-
thiophenecarboxamide

Prepared by the method of Example 9(b) using the product of Example 6 and N-(2-chloroethyl)morpholine.

MS (EI) 390 (M)⁺.

EXAMPLE 19

3-[(Aminocarbonyl)amino]-5-{3-[2-(1-pyrrolidinyl)ethoxy]phenyl}-2-thiophenecarboxamide

- 5 Prepared by the method of Example 9(b) using the product of Example 6 and N-(2-chloroethyl)pyrrolidine.

MS (EI) 374 (M)⁺.

10

EXAMPLE 20

3-[(Aminocarbonyl)amino]-5-{3-[2-(1-piperidinyl)ethoxy]phenyl}-2-thiophenecarboxamide

- Prepared by the method of Example 9(b) using the product of Example 6 and
15 N-(2-chloroethyl)piperidine.

MS (EI) 388 (M)⁺.

EXAMPLE 21

- 20 3-[(Aminocarbonyl)amino]-5-{3-[3-(dimethylamino)propoxy]phenyl}-2-thiophenecarboxamide

Prepared by the method of Example 9(b) using the product of Example 6 and N-(3-chloropropyl)dimethylamine.

25

MS (EI) 362 (M)⁺.

EXAMPLE 22

- 3-[(Aminocarbonyl)amino]-5-{2-[2-(1-morpholinyl)ethoxy]phenyl}-2-thiophenecarboxamide
- 30

Prepared by the method of Example 9(b) but using N-(2-chloroethyl)morpholine.

MS (APCI) 391 (M+H)⁺.

EXAMPLE 23

5 3-[(Aminocarbonyl)amino]-5-{2-[2-(1-pyrrolidiny)ethoxy]phenyl}-2-thiophenecarboxamide

Prepared by the method of Example 9(b) but using N-(2-chloroethyl)pyrrolidine.

MS (APCI) 375 (M+H)⁺.

10

EXAMPLE 24

3-[(Aminocarbonyl)amino]-5-{2-[2-(1-piperidiny)ethoxy]phenyl}-2-thiophenecarboxamide

15 Prepared by the method of Example 9(b) but using N-(2-chloroethyl)piperidine.

MS (APCI) 389 (M+H)⁺.

EXAMPLE 25

20 3-[(Aminocarbonyl)amino]-5-{2-[3-(dimethylamino)propoxy]phenyl}-2-thiophenecarboxamide

Prepared by the method of Example 9(b) but using N-(3-chloropropyl)dimethylamine.

MS (APCI) 363 (M+H)⁺.

EXAMPLE 26

25

2-[(Aminocarbonyl)amino]-4-methyl-5-(4-chlorophenyl)-3-thiophenecarboxamide

a) 2-Amino-4-methyl-5-(4-chlorophenyl)-3-thiophenecarboxamide

30 (4-Chlorophenyl)acetone (1.7 g), cyanoacetamide (0.84 g), sulphur (0.36 g) and morpholine (1 mL) in ethanol (5 mL) were stirred and heated at 55 °C for 6 h. The reaction mixture was cooled and filtered from a small amount of insoluble material before adding to

water (150 mL). The precipitated solid was filtered off, washed with water and then dried. The product was then triturated with ether and collected (1.0 g).

MS (EI) 266 (M)⁺.

¹H NMR (DMSO-D₆) 7.4 (2H, d), 7.3 (2H, d), 6.9 (2H, s), 6.8 (2H, s), 2.2 (3H, s).

5

b) 2-[(Aminocarbonyl)amino]-4-methyl-5-(4-chlorophenyl)-3-thiophenecarboxamide

2-Amino-4-methyl-5-(4-chlorophenyl)-3-thiophenecarboxamide (0.44 g) was suspended in acetonitrile (25 mL) and trichloroacetylisocyanate (0.2 mL) added dropwise with stirring over 10 minutes. Stirring was continued for a further 3 h at room temperature and then a 2M solution of ammonia in methanol (10 mL) was added and stirring continued for a further 2 h. The solvent was evaporated and the residue treated with water. The resultant solid was filtered off and washed with more water. The crude product was chromatographed on silica gel eluting with dichloromethane/methanol mixtures.

10

Trituration with ether gave the title urea (0.1 g).

15

MS (ES) 310 (M+H)⁺.

¹H NMR (DMSO-D₆) 10.05 (1H, s), 7.4 (2H, d), 7.35 (2H, d), 7.25 (2H, m), 6.8 (2H, s), 2.25 (3H, s).

EXAMPLE 27

20

2-[(Aminocarbonyl)amino]-4-methyl-5-(4-methylphenyl)-3-thiophenecarboxamide

Prepared by the method of Example 26 from (4-methylphenyl)acetone.

MS (ES) 290 (M+H)⁺.

¹H NMR (DMSO-D₆) 10.04 (1H, m), 7.2 (6H, m), 6.7 (2H, m), 2.3 (3H, s), 2.25 (3H, s).

25

EXAMPLE 28

2-[(Aminocarbonyl)amino]-4-ethyl-5-phenyl-3-thiophenecarboxamide

Prepared by the method of Example 26 from 1-phenyl-2-butanone.

30

MS (ES) 290 (M+H)⁺.

¹H NMR (DMSO-D₆) 9.6 (1H, m), 7.2 (7H, m), 6.6 (2H, m), 2.7 (2H, m), 1.0 (3H, t).

EXAMPLE 292-[(Aminocarbonyl)amino]-4-methyl-5-(4-methoxyphenyl)-3-thiophenecarboxamide

Prepared by the method of Example 26 from (4-methoxyphenyl)acetone.

5 MS (ES) 306 (M+H)⁺.

¹H NMR (DMSO-D₆) 10.04 (1H, s), 7.8 (1H, m), 7.25 (3H, m), 7.0 (2H, m), 6.75 (2H, m),
3.8 (3H, s), 2.2 (3H, s).

EXAMPLE 30

10 2-[(Aminocarbonyl)amino]-4-methyl-5-(4-fluorophenyl)-3-thiophenecarboxamide

Prepared by the method of Example 26 from (4-fluorophenyl)acetone.

MS (ES) 294 (M+H)⁺.

¹H NMR (DMSO-D₆) 10.05 (1H, s), 8.3 (1H, m), 7.8 (1H, m), 7.35 (2H, m), 7.2 (2H, m),
15 6.8 (2H, m), 2.2 (3H, s).

EXAMPLE 312-[(Aminocarbonyl)amino]-4-methyl-5-(3-fluorophenyl)-3-thiophenecarboxamide

20 Prepared by the method of Example 26 from (3-fluorophenyl)acetone.

MS (ES) 294 (M+H)⁺.

¹H NMR (DMSO-D₆) 10.0 (1H, s), 7.4 (3H, m), 7.2 (3H, m), 6.8 (2H, s), 2.25 (3H, s).

EXAMPLE 32

25 2-[(Aminocarbonyl)amino]-4-methyl-5-(3-methoxyphenyl)-3-thiophenecarboxamide

Prepared by the method of Example 26 from (3-methoxyphenyl)acetone.

MS (ES) 306 (M+H)⁺.

30

EXAMPLE 33

2-[(Aminocarbonyl)amino]-4-methyl-5-(3-chloro-4-methoxyphenyl)-3-
thiophenecarboxamide

Prepared by the method of Example 26 from (3-chloro-4-methoxyphenyl)acetone.

MS (ES) 340/342 (M+H)⁺.

¹H NMR (DMSO-D₆) 7.25 (5H, m), 6.8 (2H, s), 3.9 (3H, s), 2.2 (3H, s).

5

EXAMPLE 34

2-[(Aminocarbonyl)amino]-4-methyl-5-(2-chlorophenyl)-3-thiophenecarboxamide

Prepared by the method of Example 26 from (2-chlorophenyl)acetone.

10 MS (ES) 310/312 (M+H)⁺.

¹H NMR (DMSO-D₆) 10.22 (1H, s), 7.6 (1H, m), 7.4 (3H, m), 7.2 (2H, m), 6.8 (2H, s),
2.05 (3H, s).

15

EXAMPLE 35

2-[(Aminocarbonyl)amino]-4-methyl-5-(3-trifluoromethylphenyl)-3-thiophenecarboxamide

Prepared by the method of Example 26 from (3-trifluoromethylphenyl)acetone.

MS (ES) 344 (M+H)⁺.

20 ¹H NMR (DMSO-D₆) 7.65 (3H, m), 7.6 (1H, s), 7.4 (2H, m), 7.2 (2H, m), 6.8 (2H, s), 2.25
(3H, s).

EXAMPLE 36

2-[(Aminocarbonyl)amino]-4-methyl-5-(3-methyl-4-methoxyphenyl)-3-
25 thiophenecarboxamide

Prepared by the method of Example 26 from (3-methyl-4-methoxyphenyl)acetone.

MS (ES) 320 (M+H)⁺.

30 ¹H NMR (DMSO-D₆) 10.04 (1H, m), 7.2 (1H, m), 7.1 (3H, m), 6.95 (1H, d), 6.7 (2H, s),
3.8 (3H, s), 2.2 (3H, s), 2.15 (3H, s).

EXAMPLE 372-[(Aminocarbonyl)amino]-4-methyl-5-(3,5-dimethoxyphenyl)-3-thiophenecarboxamide

Prepared by the method of Example 26 from (3,5-dimethoxyphenyl)acetone.

5 MS (ES) 336 (M+H)⁺.

¹H NMR (DMSO-D₆) 6.7 (2H, m), 6.4 (3H, s), 3.8 (6H, s), 2.25 (3H, s).

EXAMPLE 382-[(Aminocarbonyl)amino]-4-methyl-5-(2,3-dimethoxyphenyl)-3-thiophenecarboxamide

10

Prepared by the method of Example 26 from (2,3-dimethoxyphenyl)acetone.

MS (ES) 336 (M+H)⁺.

¹H NMR (DMSO-D₆) 10.16 (1H, m), 7.2(1H, m), 7.05 (3H, m), 6.8 (1H, m), 6.7 (2H, m),
3.8 (3H, s), 3.5 (3H, s), 2.1(3H, s).

15

EXAMPLE 392-[(Aminocarbonyl)amino]-4-methyl-5-(4-isopropylphenyl)-3-thiophenecarboxamide

Prepared by the method of Example 26 from (4-isopropylphenyl)acetone.

20 MS (ES) 316 (M-H)⁻.

¹H NMR (DMSO-D₆) 7.25 (4H, s), 7.25 (2H, m), 6.7 (2H, m), 2.9 (1H, m), 2.25 (3H, s),
1.2 (6H, d).

EXAMPLE 40

25 2-[(Aminocarbonyl)amino]-4-methyl-5-(3,4,5-trimethoxyphenyl)-3-thiophenecarboxamide

Prepared by the method of Example 26 from (3,4,5-trimethoxyphenyl)acetone.

MS (ES) 364 (M-H)⁻.

¹H NMR (DMSO-D₆) 6.7 (2H, m), 6.6 (2H, s), 3.8 (6H, s), 3.7 (3H, s), 2.3 (3H, s).

30

EXAMPLE 412-[(Aminocarbonyl)amino]-4-methyl-5-(4-pyridyl)-3-thiophenecarboxamide

Prepared by the method of Example 26 from (4-pyridyl)acetone.

5 MS (ES) 275 (M-H).

¹H NMR (DMSO-D₆) 8.55 (2H, m), 7.2 (4H, m), 7.1 (2H, m), 2.35 (3H, s).

EXAMPLE 422-[(Aminocarbonyl)amino]-4-methyl-5-(2-pyridyl)-3-thiophenecarboxamide

10

Prepared by the method of Example 26 from (2-pyridyl)acetone.

MS (ES) 275 (M-H).

¹H NMR (DMSO-D₆) 9.9 (1H, s), 8.5 (1H, m), 7.8 (1H, m), 7.5 (1H, m), 7.4 (2H, m), 7.2
15 (2H, m), 6.7 (2H, m).

a) (2-Pyridyl)acetone

2-Picoline (2 g) was dissolved in tetrahydrofuran (30 ml) and the solution was cooled to
-75 °C. Butyl lithium (14.73 ml. of a 1.6M solution in hexane) was added dropwise and the
20 mixture stirred for 2 h. Dimethylacetamide (1.87 ml) was then added dropwise and the
reaction was allowed to warm up to room temperature and stirring was continued for a
further 2 h. Water (8.6 ml) and 36% hydrochloric acid (1.3 ml) were added and after
stirring for another 30 minutes, ethyl acetate was added. The separated solvent phase was
washed with brine and then dried (MgSO₄). On evaporation an oil was obtained and used
25 without further purification.

MS (ES) 134 (M-H).

¹H NMR (CDCl₃) 8.6 (1H, m), 7.6 (1H, m), 7.2 (2H, m), 3.9 (2H, s), 2.2 (3H, s).

EXAMPLE 432-[(Aminocarbonyl)amino]- 5-[2-(5-methoxypyridyl)]-4-methyl-3-thiophenecarboxamide

30

Prepared by the method of Example 26 from [2-(5-methoxypyridyl)]acetone.

MS (ES) 307 (M-H)⁻.

¹H NMR (DMSO-D₆) 9.93 (1H, s), 8.26 (1H, d), 7.46 - 7.37 (2H, m), 7.33 (2H, bs), 6.72 (2H, bs), 3.83 (3H, s), 2.40 (3H, s).

5 a) [2-(5-Methoxypyridyl)]acetone

Prepared in a similar manner to Example 42(a) from 3-methoxy-6-methylpyridine.

MS (ES) 166 (M+H)⁺.

¹H NMR (CDCl₃) 8.25 (1H, d), 7.22 - 7.10 (2H, m), 3.85 (5H, s), 2.22 (3H, s).

10 b) 3-Methoxy-6-methylpyridine

A solution of 3-hydroxy-6-methylpyridine (2.5 g), sodium methoxide (1.36 g) and phenyltrimethylammonium chloride (4.33 g) in dry dimethylformamide (25 ml) was heated at reflux under argon for 2.5 h. The mixture was allowed to cool then stirred at room temperature overnight. Insoluble material was removed by filtration and washed with ethanol. The filtrate was acidified with 6M hydrochloric acid and the solvent was removed *in vacuo*. The residue was then diluted with water, basified with 2M sodium hydroxide and extracted with ether. The combined extracts were washed with brine, dried (MgSO₄), filtered and evaporated. The crude product was purified by column chromatography on silica gel eluting with 3% ethyl acetate in hexane (1.55 g, 55%).

MS (ES) 124 (M+H)⁺.

¹H NMR (CDCl₃) 8.19 (1H, d), 7.10 (1H, dd), 7.05 (1H, d), 3.83 (3H, s), 2.48 (3H, s).

EXAMPLE 44

25 2-[(Aminocarbonyl)amino]-4-methyl-5-(4-pyrimidyl)-3-thiophenecarboxamide

Prepared by the method of Example 26 from (4-pyrimidyl)acetone.

MS (ES) 278 (M-H)⁻.

¹H NMR (DMSO-D₆) 9.95 (1H, s), 9.00 (1H, s), 8.64 (1H, d), 7.55 (1H, d), 7.50 (2H, bs), 6.84 (2H, bs), 2.54 (3H, s).

a) (4-Pyrimidyl)acetone

4-Methylpyrimidine (2 g) was stirred in dry tetrahydrofuran (65 ml) under argon and the solution was cooled to -78 °C. Lithium diisopropylamide (13.8 ml, 2M solution) was
5 added dropwise over 20 minutes and stirring was continued at -78 °C for 1.5 h before dropwise addition of N-methoxy-N-methylacetamide (2.49 ml). The reaction mixture was stirred at -78 °C for a further 40 minutes before allowing to warm to room temperature over 1.25 h, then partitioned between saturated aqueous sodium carbonate and ethyl acetate. The layers were separated and the aqueous phase further extracted with ethyl acetate. The
10 combined extracts were washed with brine, dried (MgSO₄), filtered and evaporated. The residue was purified by column chromatography on silica gel eluting with 40 - 50 % ethyl acetate in hexane to give an oil which crystallised on standing (0.70 g, 24%).

MS (ES) 135 (M-H)⁻.

¹H NMR (CDCl₃) 14.40 (1H, s), 8.75 (1H, s), 8.35 (1H, d), 6.74 (1H, dd), 5.29 (1H, s),
15 2.06 (3H, s).

EXAMPLE 452-[(Aminocarbonyl)amino]-4-methyl-5-(2-pyrazinyl)-3-thiophenecarboxamide

20 Prepared by the method of Example 26 from (2-pyrazinyl)acetone.

MS (ES) 278 (M+H)⁺.

¹H NMR (DMSO-D₆) 9.95 (1H, s), 8.76 (1H, d), 8.57 (1H, t), 8.42 (1H, d), 7.45 (2H, bs),
6.91 (2H, bs).

25 a) (2-Pyrazinyl)acetone

Prepared by the method of Example 44(a) from 2-methylpyrazine.

MS (ES) 135 (M-H)⁻.

¹H NMR (CDCl₃) 8.56 - 8.51 (2H, m), 8.48 (1H, d), 3.95 (2H, s), 2.28 (3H, s).

EXAMPLE 46

2-[(Aminocarbonyl)amino]-4-methyl-5-(3,4-dichlorophenyl)-3-thiophenecarboxamide

Prepared by the method of Example 26 from (3,4-dichlorophenyl)acetone.

5 MS (ES) 342 (M-H)⁺.

¹H NMR (DMSO-D₆) 10.0 (1H, s), 8.3 (2H, m), 7.6 (1H, m), 7.35 (3H, m), 6.8 (2H, m),
2.2 (3H, s).

EXAMPLE 47

10 2-[(Aminocarbonyl)amino]-4-methyl-5-(4-cyanophenyl)-3-thiophenecarboxamide

Prepared by the method of Example 26 from (4-cyanophenyl)acetone.

MS (ES) 299 (M-H)⁺.

15

EXAMPLE 48

2-[(Aminocarbonyl)amino]-4-methyl-5-(4-hydroxyphenyl)-3-thiophenecarboxamide

Prepared by demethylating 2-[(aminocarbonyl)amino]-4-methyl-5-(4-methoxyphenyl)-3-thiophenecarboxamide using boron tribromide as in Example 9(a).

20 MS (ES) 290 (M-H)⁺.

¹H NMR (DMSO-D₆) 10.02 (1H, s), 7.8 (1H, m), 7.2 (3H, m), 7.15 (2H, m), 6.8 (2H, m),
2.2 (3H, s).

EXAMPLE 49

25 2-[(Aminocarbonyl)amino]-4-methyl-5-(4-[2-(1-piperidinyl)ethoxy]phenyl)-3-thiophenecarboxamide

Prepared by the method of Example 26 using (4-[2-(1-piperidinyl)ethoxy]phenyl)acetone.

MS (ES) 401 (M-H)⁺.

¹H NMR (DMSO-D₆) 10.04(1H,s), 7.25 (3H, m), 7.1 (2H, m), 6.7 (2H, m), 4.05 (2H, m), 2.6 (2H, m), 2.4 (4H, m), 2.2 (3H, s) 1.5 (4H, m), 1.4 (2H, m).

(4-[2-(1-Piperidinyloxy]phenyl)acetone was prepared as follows:-

5

(4-Hydroxyphenyl)acetone (1.5 g), N-chloroethylpiperidine hydrochloride (2.2 g) and potassium carbonate (3.0 g) in dimethylformamide (15 mL) were stirred and heated at 80 °C for 8 h. The reaction mixture was cooled and partitioned between ethyl acetate and water. The separated solvent phase was washed twice with saturated brine and then dried
10 (MgSO₄). The resulting oil was used without further purification.
MS (ES) 262 (M+H)⁺.

EXAMPLE 50

2-[(Aminocarbonyl)amino]-4-methyl-5-(4-[2-(diethylamino)ethoxy]phenyl)-3-
15 thiophenecarboxamide

Prepared by the method of Example 26 using (4-[2-(diethylamino)ethoxy]phenyl)acetone.

¹H NMR (DMSO-D₆) 7.35 (3H, m), 7.15 (1H, m), 7.0 (2H, m), 6.8 (2H, m), 4.05 (2H, m),
20 2.8 (2H, m), 2.45 (4H, m), 2.4 (3H, s) 1.0 (6H, t).

(4-[2-(Diethylamino)ethoxy]phenyl)acetone was prepared in a similar manner to Example 49(a).

MS (ES) 249 (M+H)⁺.

25

EXAMPLE 51

2-[Aminocarbonyl)amino]-4-methyl-5-(2-furyl)-3-thiophenecarboxamide

Prepared by the method of Example 26 using 1-(2-furyl)-propan-2-one.

30

1-(2-Furyl)-propan-2-one was prepared as follows:-

a) 1-(2-Furyl)-propan-2-ol

To a solution of furan (7.93 g) in tetrahydrofuran (100 ml) cooled to 5 °C was added dropwise n-butyl lithium (80.2 ml, 1.6M in hexanes). The mixture was stirred for 2 h. A solution of propylene oxide (12.2 ml) was added dropwise and the resulting mixture was stirred at 0 °C for 1 h. The reaction mixture was quenched with saturated ammonium chloride and extracted with diethyl ether. The organics were dried (MgSO₄) and concentrated. The crude oil was distilled to give 1-(2-furyl)-propan-2-ol (3.85 g, b.p. 68-70 °C at 6.0 mm).

¹H NMR (CDCl₃) 7.35 (1H, d), 6.3 (1H, m), 6.1 (1H, d), 4.1 (1H, m), 2.7-2.9 (2H, m), 1.8 (1H, s), 1.2 (3H, d).

b) 1-(2-Furyl)-propan-2-one

To a solution of 1-(2-furyl)-propan-2-ol (3.25 g) in dichloromethane (200 ml) was added in one portion pyridinium chlorochromate (13.0 g). The resulting mixture was stirred at room temperature for 5 h and then filtered through a small bed of silica. The organics were evaporated to give the crude product which was used without further purification (3.53 g).

¹H NMR (CDCl₃) 7.4 (1H, d), 6.35 (1H, m), 6.2 (1H, d), 3.7 (2H, s), 2.2 (3H, s).

EXAMPLE 52

2-[(Aminocarbonyl)amino]-4-trifluoromethyl-5-phenyl-3-thiophenecarboxamide

a) 2-Amino-4-trifluoromethyl-5-phenyl-3-thiophenenitrile

A solution of 3,3,3-trifluoro-1-phenylpropan-2-one (1 g), malononitrile (0.39 g), sulphur (0.25 g), triethylamine (0.22 g), in ethanol (5 ml) was stirred and heated at 85 °C for 12 h. The reaction mixture was added to water (200 ml) and extracted twice into ethyl acetate (100 ml). The mixture was separated and the organic layer dried (anhydrous sodium sulfate), filtered and concentrated. The residue was chromatographed on silica gel eluting with ethyl acetate/isohexane mixtures. The solvent was removed and the product collected (0.5 g).

MS (ES) 267 (M-H)⁺.

¹H NMR (DMSO-D₆) 7.65 (2H, s), 7.35-7.45 (5H, m).

b) 2-Amino-4-trifluoromethyl-5-phenyl-3-thiophenecarboxamide

- 5 A mixture of 2-amino-4-trifluoromethyl-5-phenyl-3-thiophenenitrile (0.12 g) and concentrated sulphuric acid (1.5 ml) was stirred and heated at 50 °C for 8 h. The reaction mixture was added to saturated aqueous sodium bicarbonate until a pH of 7 was obtained. The product was extracted into ethyl acetate (100 ml) and the organic layer was dried with anhydrous sodium sulfate (3 g), filtered and concentrated. The residue was
- 10 chromatographed on silica gel eluting with ethyl acetate/isohehexane mixtures. The solvent was removed and the product collected (0.07 g).

MS (ES) 285 (M-H)⁺.

¹H NMR (DMSO-D₆) 7.35-7.45 (5H, m), 7.2 (2H, s), 6.2 (2H, s).

15 c) 2-[(Aminocarbonyl)amino]-4-trifluoromethyl-5-phenyl-3-thiophenecarboxamide

- 2-Amino-4-trifluoromethyl-5-phenyl-3-thiophenecarboxamide (0.35 g) was suspended in tetrahydrofuran (10 ml) and trichloroacetylisocyanate (0.19 g) was added dropwise with stirring over 5 minutes. Stirring was continued for 1 h at room temperature and then a 2M solution of ammonia in methanol (10 ml) was added and stirring continued for a further
- 20 12 h. A precipitate formed and was filtered off and washed with ethyl acetate (5 ml) to give the title urea (0.12 g).

MS (ES) 328 (M-H)⁺.

¹H NMR (DMSO-D₆) 9.2 (1H, s), 7.6 (2H, s), 7.35-7.45 (5H, m), 6.6 (2H, s).

25 EXAMPLE 53

2-[(Aminocarbonyl)amino]-4-methyl-5-(2-(4-methylthiazolyl))-3-thiophenecarboxamide

- Prepared by the method of Example 26(b) using 2-amino-4-methyl-5-(2-(4-
- 30 methylthiazolyl))-3-thiophenecarboxamide.

NMR (DMSO-D6) 9.9 (1H,bs), 7.45 (2H,bs), 7.19 (1H, s), 6.85 (2H, bs), 2.49 (3H, s obscured by DMSO), 2.35 (3H, s).

MS (M+H)⁺ 297.3.

5 The preparation of the starting material was achieved as follows:

a) 2-Amino-4-methyl-5-(2-(4-methylthiazolyl))-3-thiophenecarboxamide was prepared by the method of Example 52(b) from 2-amino-3-cyano-4-methyl-5-(2-(4-methylthiazolyl))thiophene.

10 NMR (DMSO-D6) 7.12 (2H,s), 7.08 (1H, s), 6.97 (2H, bs), 3.27 (3H,s), 2.44 (3H, s)
MS (M+H)⁺ 254.

b) 2-Amino-3-cyano-4-methyl-5-(2-(4-methylthiazolyl))thiophene was prepared by the method of Example 52(a) using 1-(4-methylthiazol-2-yl)-propan-2-one.

15 NMR (DMSO-D6) 7.63 (2H, s), 7.15 (1H,s), 2.26 (3H, s), 2.24 (3H,s)
MS (M+H)⁺ 236.

c) 1-(4-methylthiazol-2-yl)-propan-2-one

To a solution of 2,4 dimethylthiazole (2 g) in dry tetrahydrofuran (20 ml) at -70 °C under
20 argon was added 1.6M n-butyllithium in hexanes (12 ml) dropwise, keeping the
temperature below -70 °C. After stirring at -60 °C for 30 minutes, N-methoxy-N-
methylacetamide (1.9 ml) was added. The mixture was allowed to warm to ambient
temperature and was then partitioned between water and ethyl acetate The organic phase
was dried (MgSO₄) and the solvent removed under reduced pressure to yield a yellow oil.

25 This was purified by column chromatography using a isohexane to 40% ethyl
acetate/isohexane gradient as the eluent to yield the product as a yellow oil (630 mg, 23%).

NMR (CDCl₃) 6.84 (1H, s), 4.1 (2H, s), 2.44 (3H, s), 2.27 (3H, s).

MS (M+H)⁺ 156.

EXAMPLE 542-[(Aminocarbonyl)amino]-4-methyl-5-phenyl-3-thiophenecarboxamide5 a) 2-Cyano-3-benzyl-but-2-enoic acid amide (E/Z mixture)

A mixture of (phenyl)acetone (5 g), cyanoacetamide (3.15 g), ammonium acetate (0.29 g) and acetic acid (0.45 mL) was refluxed in toluene (50 mL) using a Dean and Stark head to remove water for 6 h. The mixture was cooled and the crystalline product was filtered off (3 g) and used without further purification.

10 MS (ES) 201 (M+H)⁺.b) 2-Amino-4-methyl-5-phenyl-3-thiophenecarboxamide

A mixture of 2-cyano-3-benzyl-but-2-enoic acid amide (E/Z mixture) (1.0 g), morpholine (0.5 mL) and sulphur (0.18 g) in ethanol (10 mL) was heated and stirred at 40 °C for 3 h.

15

After cooling, the mixture was filtered from a trace of insoluble material and the filtrate added to water. The resulting precipitate was filtered off and washed with more water, then crystallised from 2-propanol (0.35 g).

MS (ES) 233 (M+H)⁺.¹H NMR (DMSO-D₆) 7.4 (2H, m), 7.25 (3H, m), 6.9 (2H, s), 6.8 (2H, s), 2.2 (3H, s).

20

c) 2-[(Aminocarbonyl)amino]-4-methyl-5-phenyl-3-thiophenecarboxamide

To a mixture of 2-amino-4-methyl-5-phenyl-3-thiophenecarboxamide (0.18 g) in glacial acetic acid (5 mL) and water (0.5 mL) was added sodium isocyanate (101 mg). The resulting solution was stirred at room temperature for 4 h and then poured into water. The precipitate was filtered off and washed with more water. The product was

25

chromatographed on silica gel eluting with dichloromethane/methanol mixtures to give the title product as a solid (30 mg).

MS (ES) 276 (M+H)⁺.¹H NMR (DMSO-D₆) 10.05 (1H, s), 7.4 (5H, m), 7.35 (1H, m), 6.6 (2H, s), 6.4 (2H, m),

30 2.4 (3H, s).

EXAMPLE 552-[(Aminocarbonyl)amino]-4-methyl-5-(3-methyl-isoxazol-5-yl)-3-thiophenecarboxamide

Prepared by the method of Example 54 from 1-(3-methyl-isoxazol-5-yl)-propan-2-one.

5 MS (ES) 281 (M+H)⁺.

¹H NMR (DMSO-D6) 9.95 (1H, s), 7.5 (2H, bs), 6.9 (2H, bs), 6.4 (1H, s), 2.4 (3H, s), 2.25 (3H, s).

The starting 1-(3-methyl-isoxazol-5-yl)-propan-2-one was prepared as follows:

10 To a solution of 3,5-dimethylisoxazole (5.28 g) in tetrahydrofuran (80 ml), cooled to -75 °C, was added dropwise n-butyl lithium (37.4 ml, 1.6M solution in hexanes). After completion of the addition the mixture was stirred at -75 °C for 2 h. A solution of N-methoxy-N-methylacetamide in tetrahydrofuran (10 ml) was added dropwise over 15 minutes. The mixture was allowed to warm to room temperature and then to stir for a further 2 h. The mixture was quenched with saturated ammonium acetate and extracted with diethyl ether. The organics were combined, dried (MgSO₄) and concentrated. The crude product was chromatographed on silica gel eluting with a 1:1 mixture of diethyl ether/hexane to give the title compound as an oil (1.57 g).

MS (ES) 140 (M+H)⁺.

20 ¹H NMR (CDCl₃) 6.1 (1H, s), 3.8 (2H, s), 2.3 (3H, s), 2.2 (3H, s).

EXAMPLE 562-[(Aminocarbonyl)amino]-5-(4-cyanophenyl)-3-thiophenecarboxamide

25 a) 2-Amino-3-cyanothiophene.

2,5-Dihydroxy-1,4-dithiane (14.3 g) was suspended in ethanol (250 ml) and malononitrile (13.0 g) added. The mixture was cooled to 5 °C and diethylamine (20.6 ml) in ethanol (15 ml) was added at a rate such that the temperature was maintained at 5 °C. The mixture was then heated at 30-35 °C for 1.5 h. Water (280 ml) was added and the mixture poured

onto crushed ice (400 g). After a short period of time pale brown crystals formed which were filtered off and dried on the filter (14.6 g).

MS (ES) 125 (M+H)⁺.

¹H NMR (CDCl₃) 6.7 (1H, d), 6.4 (1H, d), 4.8 (2H, bs).

5

b) 2-Acetylamino-3-cyanothiophene

2-Amino-3-cyanothiophene (12 g) was heated at reflux in acetic anhydride (34 ml) for 15 minutes, cooled and refrigerated for 3 h. The crystalline product was filtered off (13.6 g).

MS (ES) 167 (M+H)⁺.

10 ¹H NMR (DMSO-D₆) 11.6 (1H, bs), 7.1 (2H, m), 2.1 (3H, s).

c) 2-Acetylamino-5-bromo-3-cyanothiophene

2-Acetylamino-3-cyanothiophene (13.5 g) was dissolved in dimethylformamide (110 ml) and cooled in an ice/water bath. *N*-Bromosuccinimide (15.9 g) was added portion wise over 20 minutes and then the mixture warmed to room temperature and stirred for 3 h. The mixture was concentrated to approximately half the volume and water added to precipitate the product. This was filtered off and dried at 60 °C under vacuum (18.8 g).

15 ¹H NMR (DMSO-D₆) 12.0 (1H, bs), 7.4 (1H, s), 2.1 (3H, s).

20 d) 2-Acetylamino-3-cyano-5-(4-cyanophenyl)thiophene

2-Acetylamino-5-bromo-3-cyanothiophene (500 mg), 4-cyanophenylboronic acid (360 mg) and potassium carbonate (845 mg) were added to dimethoxyethane (15 ml) and water (2 ml) and the system purged with argon for 15 minutes.

Tetrakis(triphenylphosphine)palladium(0) (236 mg) was added and the mixture heated at 25 80 °C for 3.25 h. The mixture was cooled, concentrated under reduced pressure to remove most of the dimethoxymethane, dichloromethane added and the mixture filtered to give the product as a pale brown solid (470 mg).

MS (ES) 266 (M-H)⁻.

¹H NMR (DMSO-D₆) 7.8 (5H, m), 2.1 (3H, s).

30

e) 2-Amino-5-(4-cyanophenyl)-3-thiophenecarboxamide

2-Acetylamino-3-cyano-5-(4-cyanophenyl)thiophene (470 mg) was heated at reflux in ethanol (15 ml) and concentrated sulphuric acid (1.5 ml) for 2.5 h. The reaction mixture was cooled and concentrated under reduced pressure. The residue was basified with

5 MS (ES) 242 (M-H)⁺.

¹H NMR (DMSO-D₆) 7.8 (1H, s), 7.7 (4H, m), 7.5 (2H, d), 7.3 (1H, bs), 7.0 (1H, bs).

f) 2-[(Aminocarbonyl)amino]-5-(4-cyanophenyl)-3-thiophenecarboxamide

Prepared by the method of Example 26(b).

10 MS (ES) 285 (M-H)⁺.

¹H NMR (DMSO-D₆) 11.0 (1H, bs), 8.0 (1H, s), 7.8 (2H, d), 7.7 (3H, m), 7.4 (1H, bs), 7.0 (2H, bs).

EXAMPLE 57

15 2-[(Aminocarbonyl)amino]-5-(4-trifluoromethylphenyl)-3-thiophenecarboxamide

Prepared by the methods of Example 56(d-f) but using 4-trifluoromethylphenylboronic acid.

MS (ES) 328 (M-H)⁺.

20 ¹H NMR (DMSO-D₆) 11.0 (1H, bs), 7.9 (1H, s), 7.7 (5H, m), 7.3 (1H, bs), 7.0 (2H, bs).

EXAMPLE 58

2-[(Aminocarbonyl)amino]-5-(2,4-difluorophenyl)-3-thiophenecarboxamide

25

Prepared by the method of Example 56(d-f) but using 2,4-difluorophenylboronic acid.

MS (ES) 296 (M-H)⁺.

¹H NMR (DMSO-D₆) 11.0 (1H, bs), 7.7 (2H, m), 7.6 (1H, m), 7.3 (2H, m), 7.2 (1H, m), 7.0 (2H, bs).

30

EXAMPLE 592-[(Aminocarbonyl)amino]-5-(2-pyridyl)-3-thiophenecarboxamide

Prepared by the method of Example 26(b) from 2-amino-5-(2-pyridyl)-3-thiophenecarboxamide.

MS (ES) 263 (M+H)⁺.

¹H NMR (DMSO-D₆) 11.04 (1H, s), 8.46 - 8.41 (1H, m), 7.99 (1H, s), 7.81 - 7.73 (1H, m), 7.65 (1H, bs), 7.61 (1H, d), 7.27 (1H, bs), 7.19 - 7.12 (1H, m), 6.95 (2H, bs).

The starting material was prepared as follows:

a) 2-(2-Methoxyvinyl)pyridine

A stirred suspension of methoxymethyltriphenyl phosphonium chloride (12.48 g) in tetrahydrofuran (60 ml) under argon was cooled in an ice-bath. Potassium *tert*-butoxide (36.41 ml, 1M solution in tetrahydrofuran) was then added dropwise over 30 minutes to give a deep orange - red colour. Stirring was continued at 0 – 5 °C for 50 minutes then the mixture was cooled to - 78 °C. 2-Pyridinecarboxaldehyde was added dropwise and stirring continued at - 78 °C for a further 2 h then allowed to warm to room temperature and stirred for 2 h. Hexane (100 ml) was added, the mixture filtered and the filtrate evaporated *in vacuo*. The residue was purified by column chromatography on silica gel eluting with 10% ethyl acetate in hexane to give the pure *cis* -2-(2-methoxyvinyl)pyridine (0.91 g, 24%):

¹H NMR (CDCl₃) 8.51 (1H, d), 7.88 (1H, d), 7.63 - 7.55 (1H, m), 7.02 (1H, dd), 6.35 (1H, d, *J* = 7 Hz), 5.50 (1H, d, *J* = 7 Hz), 3.84 (3H, s);

and a mixture of *cis* : *trans* products (1 : 1, 2.54 g, 67%):

¹H NMR (CDCl₃) 8.52 - 8.49 (0.5H, m), 8.46 - 8.41 (0.5H, m), 7.86 (0.5H, d), 7.63 - 7.48 (1.5H, m), 7.07 - 6.95 (1.5H, m), 6.35 (0.5H, d, *J* = 7 Hz), 5.87 (0.5H, d, *J* = 13 Hz), 5.50 (0.5H, d, *J* = 7 Hz), 3.84 (1.5H, s), 3.73 (1.5H, s).

b) 2-Amino-5-(2-pyridyl)-3-thiophenecarboxamide

2-(2-Methoxyvinyl)pyridine (1.28 g) was dissolved in ethanol (13 ml) and to the solution was added 6M sulphuric acid (3.6 ml). The solution was heated to 80 °C for 20 minutes then allowed to cool to 55 °C. Morpholine (8 ml) was added followed by cyanoacetamide (0.796 g) and sulfur (0.334 g). The mixture was heated at 55 °C for 4 h. After cooling to
5 room temperature the solution was poured into water (100 ml) and extracted with ethyl acetate. The extracts were dried (MgSO₄), filtered and evaporated. The residue was adsorbed onto silica gel and eluted with 2 - 5% methanol in dichloromethane to give an orange solid (345 mg, 17%).

MS (ES) 220 (M+H)⁺.

10 ¹H NMR (DMSO-D₆) 8.36 (1H, dd), 7.83 (1H, s), 7.76 - 7.67 (1H, m), 7.62 (2H, s), 7.54 (1H, d), 7.25 (1H, bs), 7.12 - 7.05 (1H, m), 6.83 (1H, bs).

EXAMPLE 60

15 2-[(Aminocarbonyl)amino]-5-(3-pyridyl)-3-thiophenecarboxamide

Prepared by the method of Example 59 from 3-(2-methoxyvinyl)pyridine.

MS (ES) 263 (M+H)⁺.

¹H NMR (DMSO-D₆) 11.02 (1H, s), 8.74 (1H, d), 8.41 (1H, dd), 7.84 (1H, dd), 7.82 (1H, s), 7.66 (1H, bs), 7.38 (1H, dd), 7.33 (1H, bs), 6.98 (2H, bs).

20

3-(2-Methoxyvinyl)pyridine

Prepared by the method of Example 59(a) from 3-pyridinecarboxaldehyde.

1 : 2.1 *cis* : *trans* products.

25 ¹H NMR (CDCl₃) 8.66 (0.32H, d), 8.47 (0.68H, d), 8.39 - 8.32 (1H, m), 7.99 - 7.92 (0.32H, m), 7.55 - 7.49 (0.68H, m), 7.21 - 7.12 (1H, m), 7.06 (0.68H, d, *J* = 13 Hz), 6.25 (0.32H, d, *J* = 7 Hz), 5.75 (0.68H, d, *J* = 13 Hz), 5.20 (0.32H, d, *J* = 7 Hz), 3.80 (0.96H, s), 3.70 (2.04H, s).

EXAMPLE 61

30 2-[(Aminocarbonyl)amino]-5-(4-pyridyl)-3-thiophenecarboxamide

Prepared by the method of Example 59 from 4-(2-methoxyvinyl)pyridine.

MS (ES) 263 (M+H)⁺.

¹H NMR (DMSO-D6) 11.09 (1H, s), 8.50 (2H, d), 8.03 (1H, s), 7.72 (1H, bs), 7.44 (2H, d), 7.35 (1H, bs), 7.04 (2H, bs).

5

4-(2-Methoxyvinyl)pyridine

Prepared by the method of Example 59(a) from 4-pyridinecarboxaldehyde.

1 : 1.13 *cis* : *trans* products.

MS (EI) 135 (M⁺).

10 ¹H NMR (CDCl₃) 8.48 (0.94H, d), 8.43 (1.06H, d), 7.41 (0.94H, d), 7.25 (0.53H, d, *J* = 14 Hz), 7.08 (1.06H, d), 6.32 (0.47H, d, *J* = 8 Hz), 5.70 (0.53H, d, *J* = 14 Hz), 5.17 (0.47H, d, *J* = 8 Hz), 3.85 (1.41H, s), 3.73 (1.59H, s).

EXAMPLE 62

15 2-[(Aminocarbonyl)amino]-5-[5-(2-methoxypyridyl)]-3-thiophenecarboxamide

Prepared by the method of Example 59 from 2-methoxy-5-(2-methoxyvinyl)pyridine.

MS (ES) 293 (M+H)⁺.

20 ¹H NMR (DMSO-D6) 10.96 (1H, bs), 8.27 (1H, d), 7.80 (1H, dd), 7.61 (1H, s), 7.61 (1H, bs), 7.28 (1H, bs), 6.95 (2H, bs), 6.85 (1H, d), 3.84 (3H, s).

2-Methoxy-5-(2-methoxyvinyl)pyridine

Prepared by the method of Example 59(a) from 5-(2-methoxypyridine)carboxaldehyde.

25 1 : 1.44 *cis* : *trans* products.

MS (EI) 165 (M⁺).

¹H NMR (CDCl₃) 8.24 (0.41H, d), 7.98 (0.59H, d), 7.91 (0.41H, dd), 7.47 (0.59H, dd), 6.92 (0.59H, d, *J* = 13 Hz), 6.70 - 6.63 (1H, m), 6.11 (0.41H, d, *J* = 7 Hz), 5.72 (0.59H, d, *J* = 13 Hz), 5.14 (0.41H, d, *J* = 7 Hz), 3.92 and 3.90 (3H, s), 3.76 (1.23H, s), 3.68 (1.77H, s).

30

The 5-(2-methoxypyridine)carboxaldehyde was prepared as follows:

a) Bromine (0.99 ml) was added dropwise to a stirred suspension of sodium acetate (1.59 g) and 2-methoxypyridine (1.93 ml) in acetic acid (10 ml). The reaction mixture was stirred at room temperature for 25 minutes, then at 80 °C for 2.5 h. The mixture was then allowed to cool and poured into ice-water, neutralised with 2M sodium hydroxide and extracted with ether. The combined extracts were dried (MgSO₄), filtered and evaporated. The crude product was purified by column chromatography on silica gel eluting with 5% ethyl acetate in hexane to give 5-bromo-2-methoxypyridine as a colourless oil (1.75 g, 51%).

MS (ES) 190, 188 (M+H)⁺.

¹H NMR (CDCl₃) 8.20 (1H, d), 7.63 (1H, dd), 6.65 (1H, d), 3.90 (3H, s).

b) 5-Bromo-2-methoxypyridine (1.53 g) was stirred in dry tetrahydrofuran (35 ml) under argon at - 78 °C. Butyl lithium (6.6 ml, 1.6M solution) was added dropwise to the solution and stirring continued at - 78 °C for 1.5 h. Dimethylformamide (1.3 ml) was then added dropwise and stirring continued at - 78 °C for a further 30 minutes before allowing to warm to room temperature. The reaction mixture was poured into saturated aqueous sodium hydrogen carbonate and the aqueous phase was extracted with ether. The combined extracts were dried (MgSO₄), filtered and evaporated. The residue was purified by column chromatography on silica gel to give 5-(2-methoxypyridine)carboxaldehyde as a white solid (0.91 g, 81%).

¹H NMR (CDCl₃) 9.95 (1H, s), 8.63 (1H, d), 8.06 (1H, dd), 6.85 (1H, d), 4.04 (3H, s).

25

EXAMPLE 63

2-[(Aminocarbonyl)amino]-5-[5-(2,4-dimethoxypyrimidyl)]-3-thiophenecarboxamide

Prepared by the method of Example 59 from 2,4-dimethoxy-5-(2-methoxyvinyl)-pyrimidine.

MS (ES) 324 (M+H)⁺.

30

¹H NMR (DMSO-D₆) 11.01 (1H, s), 8.50 (1H, s), 7.70 (1H, s), 7.69 (1H, bs), 7.31 (1H, bs), 6.95 (2H, bs), 4.05 (3H, s), 3.94 (3H, s).

5-(2,4-Dimethoxypyrimidine)carboxaldehyde

- 5 Prepared by the method of Example 62(b) from 5-bromo-2,4-dimethoxypyrimidine.
MS (EI) 168 (M⁺).

¹H NMR (CDCl₃) 10.17 (1H, s), 8.78 (1H, s), 4.11 (3H, s), 4.09 (3H, s).

2,4-Dimethoxy-5-(2-methoxyvinyl)-pyrimidine

- 10 Prepared by the method of Example 59(a) from 5-(2,4-dimethoxypyrimidine)-
carboxaldehyde.

29% *trans* product isolated:

MS (EI) 196 (M⁺).

- ¹H NMR (CDCl₃) 8.06 (1H, s), 7.10 (1H, d, *J* = 13 Hz), 5.64 (1H, d, *J* = 13 Hz), 4.02 (3H,
15 s), 3.97 (3H, s), 3.67 (3H, s).

49% *cis* product isolated:

¹H NMR (CDCl₃) 8.95 (1H, s), 6.19 (1H, d, *J* = 7 Hz), 5.30 (1H, d, *J* = 7 Hz), 3.97 (6H, s),
3.75 (3H, s).

20

EXAMPLE 64

2-[(Aminocarbonyl)amino]-5-(4-hydroxyphenyl)-3-thiophenecarboxamide

a) 2-Amino-3-thiophenecarboxamide

- A suspension of 2,5-dihydroxy-1,4-dithiane (25 g) and cyanoacetamide (19.3 g) in ethanol
25 (120 mL) was stirred and heated to 50 °C. Triethylamine (9.2 ml) was added over 15
minutes and the mixture was stirred at 50 °C for a further 2 h. After ice cooling the solid
was filtered off and dried (21.4 g).

MS (ES) 143 (M+H)⁺.

- 30 b) 2-[(Aminocarbonyl)amino]-3-thiophenecarboxamide

Prepared by the method of Example 26(b) from 2-amino-3-thiophenecarboxamide.

MS (ES) 186 (M+H)⁺.

c) 2-[(Aminocarbonyl)amino]-5-bromo-3-thiophenecarboxamide

- 5 2-[(Aminocarbonyl)amino]-3-thiophenecarboxamide (1.0 g) was dissolved in acetic acid (20 ml) and a solution of bromine (0.35 ml) in acetic acid (5 ml) was added over 5 minutes with rapid stirring. The mixture was stirred for 90 minutes and then added to water (50 ml). The product was filtered off and washed with water and dried under vacuum (0.55 g). MS (ES) 262/264 (M-H)⁻.

- 10 ¹H NMR (DMSO-D₆) 10.63 (1H, s), 7.9 (1H, m), 7.8 (1H, s), 7.35 (1H, m), 7.15 (1H, m).

d) 2-[(Aminocarbonyl)amino]-5-(4-methoxyphenyl)-3-thiophenecarboxamide

- A solution of 2-[(aminocarbonyl)amino]-5-bromo-3-thiophenecarboxamide (0.55 g), sodium carbonate (0.44 g) and 4-methoxyphenylboronic acid (0.51 g) in dimethoxyethane
15 (60 ml) and water (2 ml) was purged with argon for 10 minutes. Tetrakis(triphenylphosphine)palladium (0.243 g) was then added and the mixture refluxed with stirring for 18 h. After cooling, the mixture was screened and evaporated. The residue was partitioned between ethyl acetate and 2N sodium hydroxide and the solid interface layer was filtered off (0.2 g).
20 MS (ES) 290 (M-H)⁺.
¹H NMR (DMSO-D₆) 10.54 (1H, s), 8.0 (1H, m), 7.9 (1H, s), 7.45 (2H, d), 7.35 (1H, m), 6.95 (2H, d), 3.8 (3H, s).

e) 2-[(Aminocarbonyl)amino]-5-(4-hydroxyphenyl)-3-thiophenecarboxamide

- 25 Prepared by the method of Example 9(a).
MS (ES) 276 (M-H)⁻.
¹H NMR (DMSO-D₆) 10.12 (1H, s), 8.0 (1H, m), 7.85 (1H, s), 7.4 (2H, d), 7.35 (1H, m), 6.9 (2H, d).

EXAMPLE 652-[(Aminocarbonyl)amino]-5-(4-chlorophenyl)-3-thiophenecarboxamide

5 Prepared by the method of Example 64(d) using 4-chlorophenylboronic acid.

MS (ES) 294 (M-H)⁺.

¹H NMR (DMSO-D₆) 10.6(1H,s), 8.1(1H,s), 7.85(1H,s), 7.5(2H,d), 7.4(3H,m), 7.0(2H,m).

EXAMPLE 66

10

2-[(Aminocarbonyl)amino]-5-(4-methanesulphonylphenyl)-3-thiophenecarboxamide

Prepared by the method of Example 64(d) using 4-methanesulphonylphenylboronic acid.

MS (ES) 338.28 (M+H)⁺.

15 ¹H NMR (DMSO-D₆) 11.06 (1H, s), 7.95 (1H, s), 7.90 (2H, d), 7.70 (3H, m), 7.35 (1H, s),

7.00 (2H, s), 3.20 (3H, s).

EXAMPLE 67

20

2-[(Aminocarbonyl)amino]-5-(2-(N-t-butoxycarbonyl)pyrrolyl)-3-thiophenecarboxamide

Prepared by the method of Example 64(d) from 1-(t-butoxycarbonyl)pyrrolyl-2-boronic acid.

MS (ES) 351 (M+H)⁺.

25 ¹H NMR (DMSO-D₆) 10.97 (1H, s), 7.55 (1H, s), 7.30 (1H, s), 7.2 (1H, s), 7.18 (1H, s),

6.85 (2H, m), 6.25 (2H, m), 1.40 (9H, s).

EXAMPLE 68

30

2-[(Aminocarbonyl)amino]-5-(2-(5-cyanothienyl))-3-thiophenecarboxamide

Prepared by the method of Example 64(d) from 5-cyanothiophenyl-2-boronic acid

MS (ES) 291 (M-H)⁺.

¹H NMR (DMSO-D₆) 11.1 (1H, s), 7.89 (1H, s), 7.85 (1H, d), 7.75 (1H, s), 7.4 (1H, s), 7.2 (1H, d), 7.1 (2H, s).

5

EXAMPLE 69

2-[(Aminocarbonyl)amino]-5-(3,5-dimethyl-isoxazol-4-yl)-3-thiophenecarboxamide

Prepared by the method of Example 64(d) from 3,5 dimethylisoxazolyl-4-boronic acid

10 MS (ES) 279 (M-H)⁺.

¹H NMR (DMSO-D₆) 11.0 (1H, s), 7.8 (1H, s), 7.4 (1H, s), 7.3 (1H, s), 6.9 (2H, s), 2.53 (3H, s), 2.3 (3H, s).

EXAMPLE 70

15

2-[(Aminocarbonyl)amino]-5-(3-furyl)-3-thiophenecarboxamide

Prepared by the method of Example 64(d) from 3-furylboronic acid.

MS (ES) 250 (M-H)⁺.

20

¹H NMR (DMSO-D₆) 10.9 (1H, s), 7.9 (1H, s), 7.7 (1H, m), 7.6 (1H, s), 7.4 (1H, s), 7.2 (1H, s), 6.9 (2H, s), 6.5 (1H, m).

EXAMPLE 71

25

2-[(Aminocarbonyl)amino]-5-(2-pyrrolyl)-3-thiophenecarboxamide

2-[(Aminocarbonyl)amino]-5-(2-(N-t-butoxycarbonyl)pyrrolyl)-3-thiophenecarboxamide (0.1 g), water (0.1 ml) and trifluoroacetic acid (2 ml) were stirred at room temperature for 8 minutes before dropwise addition to saturated aqueous sodium bicarbonate solution

(15 ml). The product was extracted into ethyl acetate and the organic layer separated. The crude product was chromatographed on silica gel eluting with methanol/dichloromethane mixtures. The solvent was removed and the product collected (0.04 g).

MS (ES) 249 (M-H)⁺.

5 ¹H NMR (DMSO-D₆) 11.04 (1H, s), 10.86 (1H, s), 7.5(1H, s), 7.2-7.15 (2H, m), 6.85 (2H, s), 6.7(1H, m), 6.15 (1H, m), 6.05 (1H, m).

EXAMPLE 72

10 2-[(Aminocarbonyl)amino]-5-(5-pyrimidinyl)-3-thiophenecarboxamide

Triisopropyl borate (1.48 ml) was added to a stirred solution of 5-bromopyrimidine (200 mg) in tetrahydrofuran (10 ml) under argon. The solution was then cooled to - 78 °C and n-butyl lithium (3.30 ml, 1.6M solution in hexanes) was added dropwise. Stirring was continued at - 78 °C for 5 minutes before allowing the reaction mixture to warm to room temperature. The solvent was removed *in vacuo*, dimethoxyethane (12 ml) was added, followed by 2-[(aminocarbonyl)amino]-5-bromo-3-thiophenecarboxamide (200 mg) and saturated aqueous sodium hydrogen carbonate (3.5 ml). The flask was purged with argon and tetrakis(triphenylphosphine) palladium (0) (90 mg) added. The mixture was heated at 90 °C for 4 h, then allowed to cool. The solvent was removed *in vacuo* and the residue taken up in 2M sodium hydroxide and 10% methanol in dichloromethane. The layers were separated and the aqueous phase was filtered to remove a small amount of insoluble material. The filtrate was then neutralised with 6M hydrochloric acid and the precipitate formed collected by filtration, washed with water and dried. The product was then trituated with methanol, collected by filtration and dried under high vacuum (47 mg, 24%).

MS (ES) 264 (M+H)⁺.

¹H NMR (DMSO-D₆) 11.02 (1H, bs), 9.01 (1H, s), 8.91 (2H, s), 7.93 (1H, s), 7.66 (1H, bs), 7.39 (1H, bs), 7.04 (2H, bs).

EXAMPLE 732-[(Aminocarbonyl)amino]-5-(2-(5-chlorothieryl))-3-thiophenecarboxamide

Prepared by the method of Example 72 using 5-chloro-2-bromothiophene.

5 MS(ES) 300.18 (M-H)⁻.

¹H NMR (DMSO-D₆) 11.0(1H, s), 7.75(1H,s), 7.50(1H,s), 7.25(1H,s), 7.0(1H,d), 6.95 (3H, d+bs).

EXAMPLE 74

10 2-[(Aminocarbonyl)amino]-5-[2-(5-trifluoromethylpyridyl)]-3-thiophenecarboxamide

Prepared in a similar manner to Example 72 from 2-bromo-5-trifluoromethylpyridine.

MS (ES) 331 (M+H)⁺.

¹H NMR (DMSO-D₆, 400 MHz) 11.17 (1H, s), 8.85 (1H, s), 8.26 (1H, s), 8.21 (1H, d),
15 7.83 (1H, d), 7.76 (1H, bs), 7.39 (1H, bs), 7.07 (2H, bs).

EXAMPLE 752-[(Aminocarbonyl)amino]-5-[2-(5-bromopyridyl)]-3-thiophenecarboxamide

20 Prepared in a similar manner to Example 72 from 2,5-dibromopyridine.

MS (ES) 343, 341 (M+H)⁺.

¹H NMR (DMSO-D₆, 500 MHz) 11.07 (1H, s), 8.55 (1H, d), 8.03 (1H, s), 8.02 (1H, dd),
7.63 (1H, bs), 7.58 (1H, d), 7.26 (1H, bs), 6.95 (2H, bs).

25

EXAMPLE 762-[(Aminocarbonyl)amino]-5-(2-(5-cyanofuryl))-3-thiophenecarboxamide

Prepared by the method of Example 72 using 5-cyano-2-bromofuran.

MS(ES) 275 (M-H)⁻.

¹H NMR (DMSO-D₆) 11.1 (1H, bs), 7.85 (1H, s), 7.8 (1H, bs), 7.6 (1H, d), 7.35 (1H, bs), 7.1 (2H, bs), 6.75 (1H, d).

EXAMPLE 77

5 2-[(Aminocarbonyl)amino]-5-(4-[2-(1-piperidinyloxy)]phenyl)-3-thiophenecarboxamide

Prepared as in Example 72 using 4-[2-(1-piperidinyloxy)]bromobenzene.

MS(ES) 389(M+H)⁺.

10 ¹H NMR (DMSO-D₆) 10.98 (1H, s), 7.62 (1H, s), 7.6 (1H, s), 7.42 (2H, d), 7.25 (1H, d), 6.98 (2H, d), 6.9 (2H, s), 4.15 (2H, m), 1.6 (4H, M), 1.42 (2H, m).

4-[2-(1-Piperidinyloxy)]bromobenzene was prepared as follows:-

15 a) 4-Bromophenol (1 g), N-(2-chloroethyl)piperidine hydrochloride (0.94 g) and potassium carbonate (1.76 g) in dimethylformamide (15 ml) were stirred and heated at 60 °C for 15 h. The reaction mixture was cooled and partitioned between ethyl acetate and water. The separated solvent phase was washed twice with 2N sodium hydroxide, once with saturated brine and then dried (MgSO₄). The resulting oil was used without further purification.

20 MS (ES) 284 (M+H)⁺.

¹H NMR (DMSO-D₆) 7.2 (2H, d), 6.9 (2H, d), 4.05 (2H, m), 2.62 (2H, t), 2.38 (4H, m), 1.48 (4H, m), 1.36 (2H, m).

EXAMPLE 78

25

2-[(Aminocarbonyl)amino]-5-(4-[2-(1-(2,2,6,6-tetramethyl)piperidinyloxy)]phenyl)-3-thiophenecarboxamide

Prepared as in Example 72 using 4-[2-(2,2,6,6-tetramethyl-1-piperidinyloxy)]bromobenzene which was prepared in a similar manner to Example 77(a).

MS(ES) 445(M+H)⁺.

5 ¹H NMR (DMSO-D₆) 7.48 (2H, d), 6.96 (2H, d), 4.22 (2H, m), 3.62 (2H, m), 1.8 (4H, m), 1.56 (2H, m), 1.42 (6H, s), 1.36 (6H, s).

EXAMPLE 79

2-[(Aminocarbonyl)amino]-5-(4-(thiazol-4-yl-methoxy)phenyl)-3-thiophenecarboxamide

10

Prepared as in Example 72 using 4-[thiazol-4-yl-methoxy]bromobenzene which was prepared in a similar manner to Example 77(a).

MS(ES) 375(M+H)⁺.

15 ¹H NMR (DMSO-D₆) 10.91 (1H, s), 9.1 (1H, s), 7.88 (1H, s), 7.82 (1H, bs), 7.75 (1H, s), 7.42 (2H, d), 7.24 (1H, bs), 7.08 (2H, d), 6.9 (1H, bs), 5.11 (2H, s).

EXAMPLE 80

2-[(Aminocarbonyl)amino]-5-(4-[2-(dimethylamino)ethoxy]phenyl)-3-thiophenecarboxamide

20

Prepared as in Example 72 using 4-[2-(dimethylamino)ethoxy]bromobenzene which was prepared in a similar manner to Example 77(a).

MS(ES) 349(M+H)⁺.

25 ¹H NMR (DMSO-D₆) 11 (1H, s), 7.65 (1H, bs), 7.6 (1H, s), 7.5 (2H, d), 7.28 (1H, bs), 7.05 (2H, d), 6.9 (2H, bs), 4.45 (2H, t), 3.5 (2H, t), 2.85 (6H, s).

EXAMPLE 812-[(Aminocarbonyl)amino]-5-(4-[2-(diethylamino)ethoxy]phenyl)-3-thiophenecarboxamide

5

Prepared as in Example 72 using 4-[2-(dimethylamino)ethoxy]bromobenzene which was prepared in a similar manner to Example 77(a).

MS(ES) 377(M+H)⁺.

¹H NMR (DMSO-D₆) 11 (1H, s), 7.65 (1H, bs), 7.6 (1H, s), 7.5 (2H, d), 7.28 (1H, bs), 7.05
10 (2H, d), 6.9 (2H, bs), 4.35 (2H, t), 3.5 (2H, t), 3.25 (4H, m), 1.2 (6H, t).

EXAMPLE 822-[(Aminocarbonyl)amino]-5-(4-[2-(1-morpholinyl)ethoxy]phenyl)-3-thiophenecarboxamide

15

Prepared as in Example 72 using 4-[2-(1-morpholinyl)ethoxy]bromobenzene which was prepared in a similar manner to Example 77(a).

MS(ES) 391 (M+H)⁺.

¹H NMR (DMSO-D₆) 10.9 (1H, s), 7.55 (1H, s), 7.5 (2H, d), 7.15 (1H, bs), 7.05 (2H, d),
20 6.55 (2H, bs), 4.4 (2H, s), 3.8 (4H, s), 3.4-2.8 (6H, bm).

EXAMPLE 832-[(Aminocarbonyl)amino]-5-(2-furyl)-3-thiophenecarboxamide

25

To a solution of furan (598 mg) in dry tetrahydrofuran (15 ml) cooled to -75 °C under argon was added dropwise n-butyl lithium (7.16 ml, 1.6M solution in hexanes). The mixture was allowed to warm to -10 °C and stirred at this temperature for 1 h. The mixture

was cooled to -60°C and triisopropyl borate (3.04 ml) was added and after the addition the mixture was allowed to warm to room temperature and stirred for a further 0.5 h. The mixture was concentrated and dimethoxyethane (12 ml), 2-[(aminocarbonyl)amino]-5-bromo-3-thiocarboxamide and saturated sodium bicarbonate (5.5 ml) were added. The mixture was purged with argon and tetrakis(triphenylphosphine) palladium (0) (150 mg) was added and then refluxed under argon atmosphere for 4 h. The mixture was concentrated and then partitioned between ethyl acetate and 2N sodium hydroxide. The organic layer was dried (MgSO_4) and evaporated. The residue was purified by column chromatography on silica gel eluting with methanol / methylene chloride mixture to give the title compound as a solid (152 mg).

MS (ES) 235 ($\text{M}-\text{NH}_2$) $^{+}$.

^1H NMR ($\text{DMSO}-d_6$) 7.7 (1H, bs), 7.65 (1H, s), 7.5 (1H, s), 7.3 (1H, bs), 7.25 (1H, bs), 7.0 (2H, bs), 6.5 (2H, dd).

15

EXAMPLE 84

2-[(Aminocarbonyl)amino]-5-(2-(5-methylfuryl))-3-thiophenecarboxamide

Prepared in a similar manner to Example 83 from 2-methylfuran.

20 MS (ES) 266 ($\text{M}+\text{H}$) $^{+}$.

^1H NMR ($\text{DMSO}-d_6$) 11.0 (1H, bs), 7.7 (1H, bs), 7.4 (1H, s), 7.2 (1H, bs), 6.95 (2H, bs), 6.35 (1H, d), 6.1 (1H, d), 2.3 (3H, s).

25

EXAMPLE 85

5-[(Aminocarbonyl)amino]-2-(3,5-dichlorophenyl)-1,3-oxazole-4-carboxamide

Prepared as in Example 26(b) from 5-amino-2-(3,5-dichlorophenyl)-1,3-oxazole-4-carboxamide.

30 NMR ($\text{DMSO}-d_6$) 7.8 (2H, s), 7.75 (1H, s), 7.54 (1H, bs), 7.43 (1H, bs), 6.81 (2H, bs)

MS (M+H)+ 315.2/317.9.

The starting material was prepared as follows:

- 5 Concentrated sulphuric acid (5 ml) was added to 5-amino-2-(3,5-dichlorophenyl)-1,3-oxazole-4-carbonitrile (490 mg) at 0 °C. After stirring at ambient temperature for 90 minutes the mixture was poured onto ice and neutralised by addition of potassium carbonate. The mixture was extracted with ethyl acetate which was then dried (MgSO₄) and evaporated under reduced pressure to yield a pale yellow solid (420 mg, 80%).
- 10 NMR (DMSO-D₆) 7.67 (2H, s), 7.63 (1H, s), 7.15 (2H, bs), 6.99 (2H, bs).
MS (M+Na)+ 294.23/296.22.

EXAMPLE 86

- 15 5-[(Aminocarbonyl)amino]-2-(4-trifluoromethylphenyl)-1,3-oxazole-4-carboxamide

Prepared as Example 26(b) from 5-amino-2-(4-(trifluoromethyl)phenyl)-1,3-oxazole-4-carboxamide to yield a cream solid (54%).

- NMR (DMSO-D₆) 9.26 (1H, bs), 8.08 (2H, d), 7.91 (2H, d), 7.52 (1H, bs), 7.43 (1H, bs),
20 6.79 (2H, bs).
MS(ES) (M+H)+ 315.28.

The starting material was made as in Example 85 but starting from 5-amino-2-(4-(trifluoromethyl)phenyl)-1,3-oxazole-4-carbonitrile to yield a cream solid (61%).

- 25 NMR (DMSO-D₆) 7.93 (2H, d), 7.82 (2H, d), 7.16 (2H, bs), 6.99 (1H, bs).
MS (M-H)⁻ 270.3.

EXAMPLE 87

2-[(Aminothiocabonyl)amino]-5-phenyl-3-thiophenecarboxamide

A solution of 2-amino-5-phenyl-3-thiophenecarboxamide (1.09 g, 5 mmol) and trimethylsilyl isothiocyanate (0.85 ml, 6 mmol) in N,N-dimethylacetamide was stirred at 75 °C for 7 days. N,N-Dimethylformamide was added until solution. The solvent was removed and the resulting slurry was chromatographed on silica gel eluting with isohexane followed by methylene chloride and diethyl ether to give the product as a yellow solid (0.49 g, 35%).

¹H NMR (DMSO-D₆, 300MHz) δ 12.59 (1H, s), 8.40 (2H, s), 7.85 (1H, s), 7.77 (1H, s), 7.53 (3H, d+s), 7.39 (2H, t), 7.25 (1H, t).

MS (ES) 278 (M+H)⁺.

Pharmacological Evaluation of Compounds

IKK2 Filter Kinase Assay

Compounds were tested for inhibition of IKK2 using a filter kinase assay. The test compounds were dissolved to 10 mM in dimethylsulphoxide (DMSO). The compounds were then diluted 1 in 40 in kinase buffer (50 mM Tris, pH 7.4 containing 0.1 mM EGTA, 0.1 mM sodium orthovanadate and 0.1% β-mercaptoethanol). 1 in 3 serial dilutions were made from this solution with 2.5% DMSO in kinase buffer. 20 µl of compound dilution was added to wells of a 96 well plate in duplicate. 20 µl 2.5% DMSO in kinase buffer instead of compound was added to control wells (0% inhibition). 20 µl 0.5 M EDTA was added instead of compound to background wells (100 % inhibition).

10 µl of a mixture of magnesium acetate, unlabelled ATP, and ³³P-labelled ATP was added to each well made such that the final concentration was 10 mM magnesium acetate, 1 µM ATP and 0.1 µCi ³³P ATP. 20 µl of a mixture of IKK2 (0.15 µg/well), 1-53 GST-IκB (0.5 µg /well) and bovine serum albumin (BSA) (8.5 ug/well) was added to each well to start the reaction. The final reaction volume was 50 µl.

The kinase reactions were incubated at 21 °C for 80 minutes and the reaction stopped by precipitating the protein by the addition of an equal volume (50 µl) of 20 % trichloroacetic acid (TCA). The precipitate was allowed to form for 10 minutes and then filtered onto a GF/C unifilter 96 well plate. Each filter was washed twice with approximately 1 ml 2 % TCA. The filter plate was dried at 30-40 °C for 60 minutes, 20 µl scintillant was added to each well and the plate sealed and radioactivity counted on a Packard Topcount microplate scintillation counter.

IKK1 Filter Kinase Assay

- 10 The selectivity of compounds was assessed by testing them for inhibition of IKK1 using a filter kinase assay. The assay conditions were identical to the IKK2 filter kinase assay except that a mixture of IKK1 (0.25 µg/well) and 1-53 GST IκB (9 µg/well) was added to each well to start the reaction.

15 Inhibition of LPS-induced TNFα production by PBMCs

The effect of test compounds on nuclear factor kappa B (NFκB) activation in cells was assessed by measuring inhibition of tumour necrosis factor alpha (TNFα) production by human peripheral blood mononuclear cells (PBMCs) stimulated by bacterial lipopolysaccharide (LPS).

20

- Human blood (250 ml), anticoagulated with heparin, was collected from healthy volunteers. Aliquots of blood (25 ml) were layered on 20 ml Lymphoprep (Nycomed) in 50 ml polypropylene centrifuge tubes. The tubes were centrifuged (Sorval RT600B) at 2,500 rpm for 30 minutes. The cloudy layer containing PBMCs was collected with a fine tipped
25 Pasteur pipette, transferred into 8 clean polypropylene centrifuge tubes (approximately 10 ml per tube) and diluted to 50 ml with phosphate-buffered saline (PBS). These tubes were centrifuged at 2,000 rpm for 8 minutes. PBS (10 ml) was added to each cell pellet and the cells were gently re-suspended. The cells were pooled in 4 centrifuge tubes, PBS was added to each tube to make the volume up to 50 ml and the tubes were centrifuged at 1,400
30 rpm for 8 minutes. The cell pellets were again re-suspended in 10 ml PBS, pooled in 2

centrifuge tubes, the volume made up to 50 ml with PBS and the tubes centrifuged at 900 rpm for 10 minutes.

5 The final cell pellets were gently re-suspended in 10 ml tissue culture medium (RPMI containing 1% heat-inactivated human serum, L-glutamine and penicillin and streptomycin), combined into 1 tube and the volume made up to 30 ml with RPMI medium. The cells were counted and the cell suspension was diluted to 2.6×10^6 cells/ml.

10 Test compounds were dissolved in DMSO to 10 mM and diluted 1 in 250 (40 μ M) with RPMI medium. The compounds were then serially diluted 1 in 3 with 0.4% DMSO in RPMI medium. Aliquots of test compound dilutions (50 μ l) were transferred to the wells of a 96-well plate. Control wells contained 0.4% DMSO in RPMI instead of compound.

15 Aliquots of the cell suspension (100 μ l) were added to each well and the plates incubated at 37°C for 30 minutes. 50 μ l of 40 μ g/ml LPS (Sigma, L-4130) was added to wells to stimulate TNF α production by the cells and the plates were incubated overnight at 37°C. RPMI medium (50 μ l) was added to negative control wells instead of LPS. The final incubation volume was 200 μ l.

20 Plates were centrifuged for 4 minutes at 1,200 rpm and supernatants were removed for measurement of TNF α concentration. Viability of the remaining cell pellet was measured using WST-1 reagent (Boehringer Mannheim, 1044807). 100 μ l RPMI medium containing 10 μ l WST-1 reagent was added to each well and the plates were incubated for 0.5 to 3 h. The absorbance at 450 nm was then measured using a 96-well plate spectrophotometer.

25

TNF α in the supernatants (freshly harvested or stored frozen at -20°C) were measured using an enzyme-linked immunosorbant assay (ELISA). The ELISA plate was prepared by coating the wells of a 96 well plate with a sheep anti-human TNF α monoclonal antibody (100 μ l of 1 μ g/ml antibody diluted in coating buffer; 0.5 M carbonate/bicarbonate buffer, pH 9.6 containing 0.2 g/l sodium azide) and incubating overnight at 4°C. Blank

30

wells were not coated. The wells were washed once with 0.1% BSA in PBS containing 0.05% Tween (PBS/Tween) then incubated for 1 h at room temperature with 1% BSA in coating buffer (200 μ l). The wells were then washed 3 times with 0.1% BSA in PBS/Tween.

5

The samples of supernatant from the PBMC incubation were diluted 1 in 3 with 1% BSA in PBS/Tween. 100 μ l aliquots of these dilutions were added to the ELISA plate. Other wells contained 100 μ l TNF α standard (10, 3.3, 1.1, 0.37, 0.12, 0.04, 0.014 and 0 ng/ml). The ELISA plate was incubated at room temperature for 2 h before the wells were washed 3 times with 0.1% BSA in PBS/Tween. A rabbit anti-human TNF α antibody (100 μ l of a 2.5 μ g/ml solution) was added to each well and the plate incubated at room temperature for 1.5 h. The wells were then washed 3 times with 0.1% BSA in PBS/Tween. Goat anti-rabbit IgG-horse radish peroxidase conjugate (ICN, 674371; 100 μ l of a 1 in 10,000 dilution) was added to each well and the plate incubated at room temperature for 1.5 h. The wells were washed 3 times with 0.1% BSA in PBS/Tween.


10
15

Peroxidase substrate was prepared by dissolving a 1 mg TMB tablet (Sigma, T-5525) in 100 μ l DMSO (100 μ l) and adding this and 36 μ l UHPO (BDH, 30559; 1 g tablet dissolved in 25 ml distilled water) to 10 ml 0.1M citrate/acetate buffer, pH6. 100 μ l substrate was added to each well and the plate incubated in the dark at room temperature for approximately 30 minutes. The reaction was stopped by adding 25 μ l 2 M sulphuric acid to each well. The absorbance at 450 nm was measured in a 96 well plate spectrophotometer.

20

CLAIMS

1. A compound of formula (I)



5

in which:

- A represents a 5-membered heteroaromatic ring containing one or two heteroatoms
10 selected independently from oxygen, nitrogen or sulfur;

- R^1 represents a phenyl group or a 5- to 7-membered heteroaromatic ring containing one to
three heteroatoms selected independently from oxygen, nitrogen or sulfur; said phenyl or
heteroaromatic ring being optionally substituted by one or more substituents selected
15 independently from halogen, cyano, nitro, $-NR^3R^4$, $-CONR^5R^6$, $-COOR^7$, $-NR^8COR^9$,
 $-SR^{10}$, $-S(O)_mR^{10}$, $-SO_2NR^5R^6$, $-NR^8SO_2R^{10}$, C_1-C_6 alkyl, trifluoromethyl, $-(CH_2)_nR^{11}$,
 $-O(CH_2)_nR^{11}$ or $-OR^{12}$;

- R^2 represents hydrogen, halogen, cyano, nitro, $-NR^{13}R^{14}$, $-CONR^{15}R^{16}$, $-COOR^{17}$,
20 $-NR^{18}COR^{19}$, $-S(O)_mR^{20}$, $-SO_2NR^{15}R^{16}$, $-NR^{18}SO_2R^{20}$, C_1-C_2 alkyl, trifluoromethyl, C_2-C_3
alkenyl, C_2-C_3 alkynyl, trifluoromethoxy, C_1-C_2 alkoxy or C_1-C_2 alkanoyl;

X represents oxygen or sulphur;

each of R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , R^9 , R^{10} and R^{12} independently represent a hydrogen atom or C_1 - C_6 alkyl;

R^{11} represents $NR^{21}R^{22}$ where R^{21} and R^{22} are independently hydrogen or C_1 - C_6 alkyl
5 optionally substituted by C_1 - C_4 alkoxy; or R^{21} and R^{22} together with the nitrogen atom to which they are attached form a 5- or 6-membered saturated ring optionally containing a further O, S or NR^{23} group where R^{23} is hydrogen or C_1 - C_6 alkyl; or R^{11} represents OR^{24} where R^{24} represents C_1 - C_6 alkyl;

10 each of R^{13} , R^{14} , R^{15} , R^{16} , R^{17} , R^{18} , R^{19} and R^{20} independently represent a hydrogen atom or C_1 - C_2 alkyl;

m represents an integer 0, 1 or 2;

15 n represents an integer 2, 3 or 4;

and optical isomers, racemates and tautomers thereof and pharmaceutically acceptable salts or solvates thereof:

20 provided that:

when A represents thiophene, furan or pyrrole, then R^1 is not 4-pyridinyl or 3-pyrazolyl;
and


when A represents oxazole, thiazole or imidazole, then R^1 is not 3-pyridinyl or
5-pyrimidyl.

25

2. A compound of formula (I), according to Claim 1, wherein X represents oxygen.

3. A compound of formula (I), according to Claim 1 or Claim 2, in which the group A
is substituted as shown below in formula (Ia), where B and D are selected from CR^2 , S, O

30 and NR^{25} , where R^2 is as defined in Claim 1 and R^{25} is hydrogen or C_1 - C_6 alkyl:



4. A compound according to any one of Claims 1 to 3 in which the ring A is thiophene.
- 5
5. A compound according to any one of Claims 1 to 4 in which R¹ represents optionally substituted phenyl.
6. A compound according to any one of Claims 1 to 5 in which R² represents H or
- 10 methyl.
7. A compound according to Claim 6 in which R² represents H.
8. A compound of formula (I), according to claim 1, selected from:
- 15 3-[(aminocarbonyl)amino]-5-phenyl-2-thiophenecarboxamide;
- 3-[(aminocarbonyl)amino]-5-(3-chlorophenyl)-2-thiophenecarboxamide;
- 3-[(aminocarbonyl)amino]-5-(4-fluorophenyl)-2-thiophenecarboxamide;
- 3-[(aminocarbonyl)amino]-5-(4-chlorophenyl)-2-thiophenecarboxamide;
- 3-[(aminocarbonyl)amino]-5-(4-isobutylphenyl)-2-thiophenecarboxamide;
- 20 3-[(aminocarbonyl)amino]-5-(2-thienyl)-2-thiophenecarboxamide;
- 3-[(aminocarbonyl)amino]-5-(4-methoxyphenyl)-2-thiophenecarboxamide;
- 3-[(aminocarbonyl)amino]-5-(3-thienyl)-2-thiophenecarboxamide;
- 3-[(aminocarbonyl)amino]-5-(3-hydroxyphenyl)-2-thiophenecarboxamide;

- 3-[(aminocarbonyl)amino]-5-(2-chlorophenyl)-2-thiophenecarboxamide;
3-[(aminocarbonyl)amino]-5-(2-methoxyphenyl)-2-thiophenecarboxamide;
3-[(aminocarbonyl)amino]-5-{2-[2-(dimethylamino)ethoxy]phenyl}-2-
thiophenecarboxamide;
5 3-[(aminocarbonyl)amino]-5-{4-[2-(dimethylamino)ethoxy]phenyl}-2-
thiophenecarboxamide;
3-[(aminocarbonyl)amino]-5-(3-methoxyphenyl)-2-thiophenecarboxamide;
2-[(aminocarbonyl)amino]-5-phenyl-3-thiophenecarboxamide;
3-[(aminocarbonyl)amino]-5-{4-[2-(1-morpholinyl)ethoxy]phenyl}-2-
10 thiophenecarboxamide;
3-[(aminocarbonyl)amino]-5-{4-[2-(1-pyrrolidiny)ethoxy]phenyl}-2-
thiophenecarboxamide;
3-[(aminocarbonyl)amino]-5-{4-[2-(1-piperidiny)ethoxy]phenyl}-2-
thiophenecarboxamide;
15 3-[(aminocarbonyl)amino]-5-{4-[3-(dimethylamino)propoxy]phenyl}-2-
thiophenecarboxamide;
3-[(aminocarbonyl)amino]-5-{3-[2-(dimethylamino)ethoxy]phenyl}-2-
thiophenecarboxamide;
3-[(aminocarbonyl)amino]-5-{3-[2-(1-morpholinyl)ethoxy]phenyl}-2-
20 thiophenecarboxamide;
3-[(aminocarbonyl)amino]-5-{3-[2-(1-pyrrolidiny)ethoxy]phenyl}-2-
thiophenecarboxamide;

- 3-[(aminocarbonyl)amino]-5-{3-[2-(1-piperidinyloxy)]phenyl}-2-thiophenecarboxamide;
- 3-[(aminocarbonyl)amino]-5-{3-[3-(dimethylamino)propoxy]phenyl}-2-thiophenecarboxamide;
- 5 3-[(aminocarbonyl)amino]-5-{2-[2-(1-morpholino)ethoxy]phenyl}-2-thiophenecarboxamide;
- 3-[(aminocarbonyl)amino]-5-{2-[2-(1-pyrrolidino)ethoxy]phenyl}-2-thiophenecarboxamide;
- 3-[(aminocarbonyl)amino]-5-{2-[2-(1-piperidino)ethoxy]phenyl}-2-
- 10 thiophenecarboxamide;
- 3-[(aminocarbonyl)amino]-5-{2-[3-(dimethylamino)propoxy]phenyl}-2-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-4-methyl-5-(4-chlorophenyl)-3-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-4-methyl-5-(4-methylphenyl)-3-thiophenecarboxamide;
- 15 2-[(aminocarbonyl)amino]-4-ethyl-5-phenyl-3-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-4-methyl-5-(4-methoxyphenyl)-3-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-4-methyl-5-(4-fluorophenyl)-3-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-4-methyl-5-(3-fluorophenyl)-3-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-4-methyl-5-(3-methoxyphenyl)-3-thiophenecarboxamide;
- 20 2-[(aminocarbonyl)amino]-4-methyl-5-(3-chloro-4-methoxyphenyl)-3-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-4-methyl-5-(2-chlorophenyl)-3-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-4-methyl-5-(3-trifluoromethylphenyl)-3-thiophenecarboxamide;

- 2-[(aminocarbonyl)amino]-4-methyl-5-(3-methyl-4-methoxyphenyl)-3-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-4-methyl-5-(3,5-dimethoxyphenyl)-3-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-4-methyl-5-(2,3-dimethoxyphenyl)-3-thiophenecarboxamide;
- 5 2-[(aminocarbonyl)amino]-4-methyl-5-(4-isopropylphenyl)-3-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-4-methyl-5-(3,4,5-trimethoxyphenyl)-3-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-4-methyl-5-(2-pyridyl)-3-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-5-[2-(5-methoxypyridyl)]-4-methyl-3-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-4-methyl-5-(4-pyrimidyl)-3-thiophenecarboxamide;
- 10 2-[(aminocarbonyl)amino]-4-methyl-5-(2-pyrazinyl)-3-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-4-methyl-5-(3,4-dichlorophenyl)-3-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-4-methyl-5-(4-cyanophenyl)-3-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-4-methyl-5-(4-hydroxyphenyl)-3-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-4-methyl-5-(4-[2-(1-piperidinyl)ethoxy]phenyl)-3-
- 15 thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-4-methyl-5-(4-[2-(diethylamino)ethoxy]phenyl)-3-thiophenecarboxamide;
- 2-[aminocarbonyl]amino]-4-methyl-5-(2-furyl)-3-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-4-trifluoromethyl-5-phenyl-3-thiophenecarboxamide;
- 20 2-[(aminocarbonyl)amino]-4-methyl-5-(2-(4-methylthiazolyl))-3-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-4-methyl-5-phenyl-3-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-4-methyl-5-(3-methyl-isoxazol-5-yl)-3-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-5-(4-cyanophenyl)-3-thiophenecarboxamide;

- 2-[(aminocarbonyl)amino]-5-(4-trifluoromethylphenyl)-3-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-5-(2,4-difluorophenyl)-3-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-5-(2-pyridyl)-3-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-5-(3-pyridyl)-3-thiophenecarboxamide;
- 5 2-[(aminocarbonyl)amino]-5-[5-(2-methoxypyridyl)]-3-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-5-[5-(2,4-dimethoxypyrimidyl)]-3-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-5-(4-hydroxyphenyl)-3-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-5-(4-chlorophenyl)-3-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-5-(4-methanesulphonylphenyl)-3-thiophenecarboxamide;
- 10 2-[(aminocarbonyl)amino]-5-(2-(N-t-butoxycarbonyl)pyrrolyl)-3-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-5-(2-(5-cyanothienyl))-3-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-5-(3,5-dimethyl-isoxazol-4-yl)-3-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-5-(3-furyl)-3-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-5-(2-pyrrolyl)-3-thiophenecarboxamide;
- 15 2-[(aminocarbonyl)amino]-5-(5-pyrimidinyl)-3-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-5-(2-(5-chlorothienyl))-3-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-5-[2-(5-trifluoromethylpyridyl)]-3-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-5-[2-(5-bromopyridyl)]-3-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-5-(2-(5-cyanofuryl))-3-thiophenecarboxamide;
- 20 2-[(aminocarbonyl)amino]-5-(4-[2-(1-piperidinyl)ethoxy]phenyl)-3-thiophenecarboxamide;
- 2-[(aminocarbonyl)amino]-5-(4-[2-(1-(2,2,6,6-tetramethyl)piperidinyl)ethoxy]phenyl)-3-thiophenecarboxamide;

2-[(aminocarbonyl)amino]-5-(4-(thiazol-4-yl-methoxy)phenyl)-3-thiophenecarboxamide;

2-[(aminocarbonyl)amino]-5-(4-[2-(dimethylamino)ethoxy]phenyl)-3-

thiophenecarboxamide;

2-[(aminocarbonyl)amino]-5-(4-[2-(diethylamino)ethoxy]phenyl)-3-

5 thiophenecarboxamide;

2-[(aminocarbonyl)amino]-5-(4-[2-(1-morpholinyl)ethoxy]phenyl)-3-

thiophenecarboxamide;

2-[(aminocarbonyl)amino]-5-(2-furyl)-3-thiophenecarboxamide;

2-[(aminocarbonyl)amino]-5-(2-(5-methylfuryl))-3-thiophenecarboxamide;

10 5-[(aminocarbonyl)amino]-2-(3,5-dichlorophenyl)-1,3-oxazole-4-carboxamide;


5-[(aminocarbonyl)amino]-2-(4-trifluoromethylphenyl)-1,3-oxazole-4-carboxamide;

2-[(aminothiocabonyl)amino]-5-phenyl-3-thiophenecarboxamide;

and pharmaceutically acceptable salts and solvates thereof.


15 9. A process for the preparation of a compound of formula (I), according to any one of Claims 1 to 8, which comprises:

(a) reaction of a compound of formula (II):



20 wherein A, R¹ and R² are as defined in Claim 1 with an isocyanate (X = O) or an isothiocyanate (X = S); or

(b) reaction of compound of formula (III) with a compound of formula (IV)


R¹-Metal

(III)

(IV)

wherein A, X, R¹ and R² are as defined in Claim 1 and LG represents a leaving group; or

- 5 (c) reaction of compound of formula (V) with a compound of formula (VI)

R¹-LG

(V)

(VI)

wherein A, X, R¹ and R² are as defined in Claim 1 and LG represents a leaving group;

- 10 and where necessary converting the resultant compound of formula (I), or another salt thereof, into a pharmaceutically acceptable salt thereof; or converting the resultant compound of formula (I) into a further compound of formula (I); and where desired converting the resultant compound of formula (I) into an optical isomer thereof.

- 15 10. A pharmaceutical composition comprising a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as claimed in any one of claims 1 to 8 in association with a pharmaceutically acceptable adjuvant, diluent or carrier.

11. A process for the preparation of a pharmaceutical composition as claimed in Claim 10 which comprises mixing a compound of formula (I), or a pharmaceutically acceptable
- 20

salt or solvate thereof, as claimed in any one of claims 1 to 8 with a pharmaceutically acceptable adjuvant, diluent or carrier.

12. A compound of formula (I), or a pharmaceutically-acceptable salt or solvate thereof, as
5 claimed in any one of claims 1 to 8 for use in therapy.

13. Use of a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as claimed in any one of claims 1 to 8 in the manufacture of a medicament for use in therapy.

10

14. Use of a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as claimed in any one of claims 1 to 8 in the manufacture of a medicament for use in the treatment or prophylaxis of diseases or conditions in which inhibition of IKK2 activity is beneficial.

15

15. Use of a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as claimed in any one of claims 1 to 8 in the manufacture of a medicament for use in the treatment or prophylaxis of inflammatory disease.

20 16. The use as claimed in Claim 15 wherein the disease is asthma.

17. The use as claimed in Claim 15 wherein the disease is rheumatoid arthritis.

18. The use as claimed in Claim 15 wherein the disease is multiple sclerosis.

25

19. The use as claimed in Claim 15 wherein the disease is chronic obstructive pulmonary disease.

20. A method of treating an IKK2 mediated disease which comprises administering to a
30 patient a therapeutically effective amount of a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as claimed in any one of claims 1 to 8.

21. A method of treating an inflammatory disease in a patient suffering from, or at risk of, said disease, which comprises administering to the patient a therapeutically effective amount of a compound of formula (I), or a pharmaceutically acceptable salt or solvate thereof, as claimed in any one of claims 1 to 8.
22. A method according to claim 21, wherein the disease is asthma.
23. A method according to claim 21, wherein the disease is rheumatoid arthritis.
24. A method according to claim 21, wherein the disease is multiple sclerosis.
25. A method according to claim 21, wherein the disease is chronic obstructive pulmonary disease.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 01/00248

A. CLASSIFICATION OF SUBJECT MATTER

C07D 333/38, C07D 409/04, C07D 413/04, C07D 417/04, C07D 263/48, A61K 31/381.

IPC7: A61K 31/395, A61P 29/00, A61P 11/06, A61P 19/02

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC7: C07D, A61K, A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

SE,DK,FI,NO classes as above

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	STN International, file CAPLUS, CAPLUS accession no. 1984:454985, document no. 101:54985, Zayed, Ezzat Mohamed et al: "Studies on 5-aminopyrazole derivatives. Synthesis of some new fused pyrazole derivatives", Monatsh. Chem. (1984), 115(4), 431-6, See CAS RN 91099-28-4 --	1,5,6,9
X	WO 9802430 A1 (PFIZER INC.), 22 January 1998 (22.01.98), see particularly claims 1, 20 and 22 --	1-25
X	EP 0853083 A1 (PFIZER INC.), 15 July 1998 (15.07.98), see particularly claims 1,11 and 13 --	1-25



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier application or patent but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

18 June 2001

Date of mailing of the international search report

25 -06- 2001

Name and mailing address of the ISA
Swedish Patent Office
Box 5055, S-102 42 STOCKHOLM
Facsimile No. +46 8 666 02 86

Authorized officer

Nebil Gecer/BS
Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SE 01/00248

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0908456 A1 (HOECHST MARION ROUSSEL DEUTSCHLAND GMBH), 14 April 1999 (14.04.99) --	1-25
A	EP 0202538 A1 (BAYER AG), 26 November 1986 (26.11.86), see claim 1 -- -----	1-9

INTERNATIONAL SEARCH REPORT

International application No.
PCT/SE01/00248

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 20-25
because they relate to subject matter not required to be searched by this Authority, namely:
see next sheet
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/SE01/00248

Claims 20-25 relate to methods of treatment of the human or animal body by surgery or by therapy/ diagnostic methods practised on the human or animal body/Rule 39.1.(iv). Nevertheless, a search has been executed for these claims. The search has been based on the alleged effects of the compounds/compositions.

INTERNATIONAL SEARCH REPORT

Information on patent family members

28/05/01

International application No.

PCT/SE 01/00248

Patent document cited in search report			Publication date	Patent family member(s)		Publication date
WO	9802430	A1	22/01/98	AU	3044197 A	09/02/98
				BR	9710352 A	17/08/99
				CA	2260213 A	22/01/98
				EP	0912548 A	06/05/99
				HR	970378 A	31/08/98
EP	0853083	A1	15/07/98	BR	9800228 A	08/09/99
				CA	2226039 A	06/07/98
				JP	10195070 A	28/07/98
				US	6048880 A	11/04/00
EP	0908456	A1	14/04/99	AU	8791298 A	22/04/99
				BR	9803880 A	02/05/00
				CA	2249542 A	06/04/99
				CN	1214339 A	21/04/99
				CZ	9803205 A	14/04/99
				DE	19744026 A	08/04/99
				HU	9802224 A	28/12/99
				JP	11302277 A	02/11/99
				PL	329042 A	12/04/99
				TR	9801996 A	00/00/00
				US	6162819 A	19/12/00
EP	0202538	A1	26/11/86	SE	0202538 T3	
				AT	39404 T	15/01/89
				AU	5721786 A	20/11/86
				BR	8602224 A	13/01/87
				CS	258481 B	16/08/88
				CS	8603569 A	15/01/88
				DE	3529247 A	20/11/86
				DE	3661493 D	00/00/00
				DK	230086 A	18/11/86
				ES	555052 A	16/02/88
				ES	8801815 A	01/05/88
				FI	862201 A	18/11/86
				GR	861269 A	12/09/86
				HU	41244 A	28/04/87
				IL	78759 D	00/00/00
				JP	61268678 A	28/11/86
				NZ	216159 A	29/09/88
				ZA	8603645 A	28/01/87